

chain nodes :

7 8 9 10 13 14 16 18 20 21 22 29 33

ring nodes :

1 2 3 4 5 6 23 24 25 26 27 28

ring/chain nodes :

17 35

chain bonds :

6-7 7-8 8-13 9-10 13-14 14-16 14-17 17-18 18-20 20-21 21-22
22-25

ring/chain bonds :

17-35

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 23-24 23-28 24-25 25-26 26-27 27-28

exact/norm bonds :

8-13 9-10 13-14 14-16 14-17 17-18 17-35 18-20 20-21

exact bonds :

6-7 7-8 21-22 22-25

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 23-24 23-28 24-25 25-26 26-27 27-28

isolated ring systems :

containing 1 : 23 :

G1:CH2, [*1]

G2:H,CH3,Et

G3:X,CF3,CCl3,CBr3,CN

Match level :

1:Atom	2:Atom	3:Atom	4:Atom	5:Atom	6:Atom	7:CLASS	8:CLASS	9:CLASS
10:CLASS	13:CLASS	14:CLASS	16:CLASS	17:CLASS	18:CLASS	20:CLASS		
21:CLASS	22:CLASS	23:Atom	24:Atom	25:Atom	26:Atom	27:Atom	28:Atom	
29:CLASS	30:Atom	33:CLASS	34:Atom	35:CLASS				

09890219

=> d his

(FILE 'HOME' ENTERED AT 12:21:47 ON 11 JUN 2007)

FILE 'REGISTRY' ENTERED AT 12:21:55 ON 11 JUN 2007

L1 STRUCTURE UPLOADED

L2 1 S L1

L3 519 S L1 FULL

FILE 'HCAPLUS' ENTERED AT 12:27:12 ON 11 JUN 2007

L4 94 S L3

L5 2 S L4 AND MATSUOKA, H?/AU

L6 92 S L4 NOT L5

L7 1 S L6 AND SATO, T?/AU

L8 91 S L6 NOT L7

L9 0 S L8 AND TAKAHASHI, T?/AU

L10 1 S L8 AND KIM, D?/AU

L11 90 S L8 NOT L10

L12 0 S L11 AND JUNG, K?/AU

L13 0 S L11 AND PARK, C?/AU

FILE 'CAOLD' ENTERED AT 12:30:36 ON 11 JUN 2007

=> s 13

L14 0 L3

=>

09890219

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal612bxx

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals
NEWS 3 JAN 16 CA/Caplus Company Name Thesaurus enhanced and reloaded
NEWS 4 JAN 16 IPC version 2007.01 thesaurus available on STN
NEWS 5 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
NEWS 6 JAN 22 CA/Caplus updated with revised CAS roles
NEWS 7 JAN 22 CA/Caplus enhanced with patent applications from India
NEWS 8 JAN 29 PHAR reloaded with new search and display fields
NEWS 9 JAN 29 CAS Registry Number crossover limit increased to 300,000 in multiple databases
NEWS 10 FEB 15 PATDPASPC enhanced with Drug Approval numbers
NEWS 11 FEB 15 RUSSIAPAT enhanced with pre-1994 records
NEWS 12 FEB 23 KOREAPAT enhanced with IPC 8 features and functionality
NEWS 13 FEB 26 MEDLINE reloaded with enhancements
NEWS 14 FEB 26 EMBASE enhanced with Clinical Trial Number field
NEWS 15 FEB 26 TOXCENTER enhanced with reloaded MEDLINE
NEWS 16 FEB 26 IFICDB/IFIPAT/IFIUDB reloaded with enhancements
NEWS 17 FEB 26 CAS Registry Number crossover limit increased from 10,000 to 300,000 in multiple databases
NEWS 18 MAR 15 WPIDS/WPIX enhanced with new FRAGHITSTR display format
NEWS 19 MAR 16 CASREACT coverage extended
NEWS 20 MAR 20 MARPAT now updated daily
NEWS 21 MAR 22 LWPI reloaded
NEWS 22 MAR 30 RDISCLOSURE reloaded with enhancements
NEWS 23 APR 02 JICST-EPLUS removed from database clusters and STN
NEWS 24 APR 30 GENBANK reloaded and enhanced with Genome Project ID field
NEWS 25 APR 30 CHEMCATS enhanced with 1.2 million new records
NEWS 26 APR 30 CA/Caplus enhanced with 1870-1889 U.S. patent records
NEWS 27 APR 30 INPADOC replaced by INPADOCDB on STN
NEWS 28 MAY 01 New CAS web site launched
NEWS 29 MAY 08 CA/Caplus Indian patent publication number format defined
NEWS 30 MAY 14 RDISCLOSURE on STN Easy enhanced with new search and display fields
NEWS 31 MAY 21 BIOSIS reloaded and enhanced with archival data
NEWS 32 MAY 21 TOXCENTER enhanced with BIOSIS reload
NEWS 33 MAY 21 CA/Caplus enhanced with additional kind codes for German patents
NEWS 34 MAY 22 CA/Caplus enhanced with IPC reclassification in Japanese patents

NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

Updated Search

09890219

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 12:21:47 ON 11 JUN 2007

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 12:21:55 ON 11 JUN 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 10 JUN 2007 HIGHEST RN 936909-28-3

DICTIONARY FILE UPDATES: 10 JUN 2007 HIGHEST RN 936909-28-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH December 2, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>

Uploading C:\Documents and Settings\brobinson1\My Documents\stnweb\Queries\4545k.str

L1 STRUCTURE UPLOADED

=> d 11

L1 HAS NO ANSWERS

L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Updated Search

09890219

Structure attributes must be viewed using STN Express query preparation.

=> s 11

SAMPLE SEARCH INITIATED 12:26:52 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 35506 TO ITERATE

5.6% PROCESSED 2000 ITERATIONS 1 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 698859 TO 721381
PROJECTED ANSWERS: 103 TO 607

L2 1 SEA SSS SAM L1

=> s 11 full

THE ESTIMATED SEARCH COST FOR FILE 'REGISTRY' IS 171.65 U.S. DOLLARS
DO YOU WANT TO CONTINUE WITH THIS REQUEST? (Y)/N or END:y
FULL SEARCH INITIATED 12:26:57 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 710231 TO ITERATE

100.0% PROCESSED 710231 ITERATIONS 519 ANSWERS
SEARCH TIME: 00.00.10

L3 519 SEA SSS FUL L1

=> file hcaplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	175.70	175.91

FILE 'HCAPLUS' ENTERED AT 12:27:12 ON 11 JUN 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 Jun 2007 VOL 146 ISS 25
FILE LAST UPDATED: 10 Jun 2007 (20070610/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 13

Updated Search

09890219

L4 94 L3

=> s l4 and matsuoka, h?/au

2553 MATSUOKA, H?/AU

L5 2 L4 AND MATSUOKA, H?/AU

=> d l5, ibib abs hitstr, 1-2

L5 ANSWER 1 OF 2 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:157810 HCAPLUS

DOCUMENT NUMBER: 136:217049

TITLE: Preparation of cyclic peptide derivatives as motilin receptor antagonists

INVENTOR(S): Matsuoka, Hiroharu; Sato, Tsutomu

PATENT ASSIGNEE(S): Chugai Seiyaku Kabushiki Kaisha, Japan

SOURCE: PCT Int. Appl., 89 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002016404	A1	20020228	WO 2001-JP7213	20010823
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2001080120	A5	20020304	AU 2001-80120	20010823
EP 1312612	A1	20030521	EP 2001-958426	20010823
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 2003191053	A1	20031009	US 2003-362574	20030224
US 7018981	B2	20060328		
PRIORITY APPLN. INFO.:			JP 2000-253950	A 20000824
			WO 2001-JP7213	W 20010823

OTHER SOURCE(S): MARPAT 136:217049

GI



Updated Search


AB The title compds. I [$T1 = (CH_2)_m$; $T2 = (CH_2)_n$; $R1$ represents optionally substituted Ph, etc.; $R2$ represents amino, etc.; $R3$ to $R6$ each represents hydrogen, Me, etc.; V, W, X, Y, Z represent carbonyl or methylene; m is an integer of 0 to 2; and n is an integer of 0 to 3] are prepared In an in vitro test for motilin receptor antagonism, (2S-(2S,12S))-2-amino-N-(2-(3-tert-butyl-4-hydroxyphenylmethyl)-1,4,8-triaza-3,7,13-trioxocyclotridecan-12-yl)-3-(4-fluorophenyl)-N-methylpropionamide showed IC_{50} of 0.52 nM.

IT 401896-13-7P 401896-15-9P 401896-22-8P
 401896-25-1P 401896-29-5P 401896-32-0P
 401896-37-5P 401896-39-7P 401896-43-3P
 401896-45-5P 401896-50-2P 401896-52-4P
 401896-63-7P 401896-64-8P 401896-65-9P
 401896-70-6P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of cyclic peptide derivs. as motilin receptor antagonists)

RN 401896-13-7 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 401896-15-9 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


09890219



RN 401896-22-8 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-3-(1,1-dimethylethyl)-N-methyl-O-(phenylmethyl)-L-tyrosyl-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 401896-25-1 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-L-lysyl-3-(1,1-dimethylethyl)-N-methyl-L-tyrosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 401896-29-5 HCAPLUS

CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl-, phenylmethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 401896-32-0 HCAPLUS

CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 401896-37-5 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl-N-methyl-, phenylmethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 401896-39-7 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 401896-43-3 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2,N6-dimethyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 401896-45-5 HCAPLUS

CN β -Alanine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2,N6-dimethyl-L-lysyl-3-(1,1-dimethylethyl)-L-tyrosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 401896-50-2 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-3-(1,1-dimethylethyl)-N-[4-oxo-4-(phenylmethoxy)butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 401896-52-4 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2-methyl-L-lysyl-N-(3-carboxypropyl)-3-(1,1-dimethylethyl)- (9CI). (CA INDEX NAME)


Absolute stereochemistry.



RN 401896-63-7 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N6-acetyl-N2-methyl-N6-[3-[(phenylmethoxy)carbonyl]amino]propyl]-L-lysyl-3-(1,1-dimethylethyl)-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 401896-64-8 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N6-acetyl-N6-(3-aminopropyl)-N2-methyl-L-lysyl-3-(1,1-dimethylethyl)- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 401896-65-9 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N6-acetyl-N2-methyl-N6-[3-[(phenylmethoxy)carbonyl]amino]propyl]-L-lysyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 401896-70-6 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N2,N5-dimethyl-N5-[(phenylmethoxy)carbonyl]-L-ornithyl-3-(1,1-dimethylethyl)-N-[4-oxo-4-(phenylmethoxy)butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 2 OF 2 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:535162 HCAPLUS

DOCUMENT NUMBER: 133:150920

TITLE: Preparation of peptides or analogs containing substituted phenethylamine moiety as motilin receptor antagonists

INVENTOR(S): Matsuoka, Hiroharu; Sato, Tsutomu; Takahashi, Tadakatsu; Kim, Dong Ick; Jung, Kyung Yun; Park, Chan Hee

PATENT ASSIGNEE(S): Chugai Seiyaku Kabushiki Kaisha, Japan

SOURCE: PCT Int. Appl., 403 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000044770	A1	20000803	WO 2000-JP444	20000128
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2359030	A1	20000803	CA 2000-2359030	20000128
EP 1149843	A1	20011031	EP 2000-901956	20000128
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
HU 200105204	A2	20020429	HU 2001-5204	20000128
JP 3715202	B2	20051109	JP 2000-596026	20000128
NO 2001003684	A	20010928	NO 2001-3684	20010726
PRIORITY APPLN. INFO.:			JP 1999-20523	A 19990128
			JP 1999-283163	A 19991004
			WO 2000-JP444	W 20000128

09890219

OTHER SOURCE(S):
GI

MARPAT 133:150920



AB Substituted phenethylamine derivs. represented by general formula (I), hydrates of the same, or pharmaceutically acceptable salts thereof [wherein Cy is a group represented by general formula Q, an optionally substituted heterocyclic group, C3-7 cycloalkyl, or phenyl; R1, R1, R1, R1 and R5 are each hydrogen, halogeno, hydroxyl, amino, trifluoromethyl or cyano, at least one of R1-R5 being halogeno, trifluoromethyl or cyano; R6 represents hydrogen, (un)substituted linear or branched C1-3 alkyl, amino, or hydroxy; R8 represents hydrogen, Me, or ethyl; R9 represents (un)substituted linear or branched C1-6 alkyl, C2-6 alkenyl, or C2-6 alkynyl, C3-7 cycloalkyl, or (un)substituted Ph; R20 represents hydrogen, or (un)substituted linear or branched C1-3 alkyl or R9 and R20 together forms C3-7 cycloalkyl; R10 represents hydrogen, (un)substituted linear or branched C1-3 alkyl; R11 represents hydrogen or (un)substituted linear or branched C1-3 alkyl, (un)substituted carbamoyl, or carboxy; R12 represents hydroxy or linear or branched C1-4 alkoxy; R13 represents hydrogen, (un)substituted linear or branched C1-6 alkyl, C2-6 alkenyl, or alkynyl, etc.; X, Y represents carbonyl or CH2; provisos are given.], which exhibit motilin receptor antagonism and being useful as drugs for preventing digestive tract movement or high level of blood motilin. Thus, 3-methyl-2-methylaminobutyric acid 2-(3-tert-butyl-4-hydroxyphenyl)-1-(2-pyridylcarbamoyl)ethylamide (preparation given) was condensed with Boc-Phe(4-F)-OH using CMPI in the presence of Et3N in THF under ice-cooling for 4 h followed by treatment of the product with CF3CO2H in CH2Cl2 gave 2-((2-amino-3-(4-fluorophenyl)propanoyl)-N-methylamino)-3-methylbutyric acid 2-(3-tert-butyl-4-hydroxyphenyl)-1-(2-pyridylcarbamoyl)ethylamide (II). II and N-Et-Phe(4-F)-N-Me-Val-N-Et-Tyr(3-tBu)-NHET showed IC50 of 0.35 and 0.17 nM, resp., for inhibiting binding of 125I-motilin to motilin receptor preparation from mucus membrane of rabbit duodenum.

IT 287205-81-6P 287205-82-7P 287205-83-8P
287205-84-9P 287205-85-0P 287205-87-2P
287205-88-3P 287205-89-4P 287205-90-7P
287205-91-8P 287205-92-9P 287205-93-0P
287205-94-1P 287205-95-2P 287205-96-3P
287205-97-4P 287205-98-5P 287205-99-6P
287206-00-2P 287206-01-3P 287206-02-4P
287206-03-5P 287206-04-6P 287206-05-7P
287206-06-8P 287206-07-9P 287206-08-0P
287206-09-1P 287206-10-4P 287206-11-5P
287206-12-6P 287206-13-7P 287206-14-8P
287206-15-9P 287206-16-0P 287206-17-1P
287206-18-2P 287206-19-3P 287206-20-6P
287206-21-7P 287206-22-8P 287206-23-9P

Updated Search

09890219

287206-24-0P 287206-25-1P 287206-26-2P
287206-27-3P 287206-28-4P 287206-29-5P
287206-30-8P 287206-31-9P 287206-32-0P
287206-33-1P 287206-34-2P 287206-35-3P
287206-36-4P 287206-37-5P 287206-38-6P
287206-39-7P 287206-40-0P 287206-41-1P
287206-42-2P 287206-43-3P 287206-44-4P
287206-45-5P 287206-46-6P 287206-47-7P
287206-49-9P 287206-50-2P 287206-51-3P
287206-52-4P 287206-53-5P 287206-55-7P
287206-56-8P 287206-58-0P 287206-59-1P
287206-60-4P 287206-61-5P 287206-62-6P
287206-63-7P 287206-64-8P 287206-65-9P
287206-66-0P 287206-67-1P 287206-68-2P
287206-69-3P 287206-70-6P 287206-71-7P
287206-72-8P 287206-73-9P 287206-74-0P
287206-75-1P 287206-76-2P 287206-77-3P
287206-78-4P 287206-79-5P 287206-80-8P
287206-81-9P 287206-82-0P 287206-83-1P
287206-84-2P 287206-85-3P 287206-86-4P
287206-87-5P 287206-88-6P 287206-89-7P
287206-90-0P 287206-91-1P 287206-92-2P
287206-93-3P 287206-94-4P 287206-95-5P
287206-96-6P 287206-97-7P 287206-98-8P
287206-99-9P 287207-00-5P 287207-01-6P
287207-02-7P 287207-03-8P 287207-04-9P
287207-05-0P 287207-06-1P 287207-07-2P
287207-08-3P 287207-09-4P 287207-10-7P
287207-11-8P 287207-12-9P 287207-13-0P
287207-14-1P 287207-15-2P 287207-16-3P
287207-17-4P 287207-18-5P 287207-19-6P
287207-20-9P 287207-21-0P 287207-22-1P
287207-23-2P 287207-24-3P 287207-25-4P
287207-26-5P 287207-27-6P 287207-28-7P
287207-29-8P 287207-30-1P 287207-31-2P
287207-32-3P 287207-33-4P 287207-34-5P
287207-35-6P 287207-38-9P 287207-39-0P
287207-41-4P 287208-31-5P 287208-73-5P
287209-74-9P 287209-77-2P 287211-41-0P
287212-51-5P 287212-55-9P 287212-56-0P
287212-57-1P 287212-58-2P 287212-59-3P
287212-60-6P 287212-61-7P 287212-62-8P
287212-63-9P 287212-64-0P 287212-65-1P
287212-66-2P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of peptides or analogs containing substituted phenethylamine moiety

as motilin receptor antagonists and drugs for preventing digestive tract movement or high level of blood motilin)


RN 287205-81-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287205-82-7 HCAPLUS

CN L-Tyrosinamide, 4-chloro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287205-83-8 HCAPLUS

CN L-Tyrosinamide, 3,4-difluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287205-84-9 HCAPLUS

CN L-Tyrosinamide, 3-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287205-85-0 HCAPLUS

CN L-Tyrosinamide, 2-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287205-87-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- α -methyl-N-(methylsulfonyl)-, mono(trifluoroacetate) (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 287205-86-1

CMF C30 H43 F N4 O6 S

Absolute stereochemistry.

Updated Search


09890219



CM 2

CRN 76-05-1


CMF C2 H F3 O2



RN 287205-88-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-methoxy-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287205-89-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-2-pyridinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219




RN 287205-90-7 HCAPLUS
CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[(aminocarbonyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



RN 287205-91-8 HCAPLUS
CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[(aminoiminomethyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287205-92-9 HCAPLUS
CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[[[(cyanoamino)(methylamino)methylene]amino]-1-[[3-(1,1-dimethylethyl)-4-

Updated Search

09890219

hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287205-93-0 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[(aminosulfonyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287205-94-1 HCAPLUS

CN Glycinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-L-tyrosyl-ψ(CH₂-NH)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287205-95-2 HCAPLUS


CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[(methylsulfonyl)amino]ethyl]-N2-methyl- (9CI)

Updated Search

09890219

(CA INDEX NAME)


Absolute stereochemistry.



RN 287205-96-3 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-3-amino-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-3-oxopropyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287205-97-4 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-(methylsulfonyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287205-98-5 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Updated Search

09890219


Absolute stereochemistry.



RN 287205-99-6 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-(methylsulfonyl)ethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-00-2 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-1-(1,4-dihydro-6-methyl-4-oxo-2-pyrimidinyl)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]ethyl]-N2-methyl-, (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-01-3 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2,5-dioxo-4-imidazolidinyl)ethyl]-N2-methyl-, (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-02-4 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1,3,4-oxadiazol-2-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-03-5 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1,2,4-oxadiazol-5-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-04-6 HCAPLUS

Updated Search

09890219

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-05-7 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1H-1,2,4-triazol-3-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-06-8 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry..

Updated Search


09890219



RN 287206-07-9 HCAPLUS

CN L-Tyrosinamide, 2-fluoro-L-tyrosyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-
N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-08-0 HCAPLUS

CN L-Tyrosinamide, 3-fluoro-L-tyrosyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-
N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-09-1 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-10-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-11-5 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-12-6 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-13-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-14-8 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-15-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-16-0 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-17-1 HCAPLUS


CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-

Updated Search

09890219

dimethylethyl)-N,N α -dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-18-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-19-3 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287206-20-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-21-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-22-8 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287206-23-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-24-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-25-1 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219




RN 287206-26-2 HCAPLUS
CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-27-3 HCAPLUS
CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-28-4 HCAPLUS
CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-29-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-30-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-31-9 HCAPLUS


CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-

Updated Search

09890219

dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-32-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-33-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-34-2 HCAPLUS


CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-

Updated Search

09890219

dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-35-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-36-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287206-37-5 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N, α -dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-38-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-39-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287206-40-0 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-41-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-42-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-43-3 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-44-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-N,3-bis(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-45-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl-N-[(methylsulfonyl)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-46-6 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-amino-3-(4-fluorophenyl)propyl]-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-47-7 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-amino-3-(4-fluorophenyl)propyl]-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-49-9 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-amino-3-(4-fluorophenyl)propyl]-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Updated Search

09890219


Absolute stereochemistry.



RN 287206-50-2 HCAPLUS

CN Butanamide, 2-[[(2S)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-51-3 HCAPLUS

CN Butanamide, 2-[[(2S)-2-amino-3-(4-fluorophenyl)propyl]methylamino]-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-52-4 HCAPLUS

CN Butanamide, 2-[[(2S)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]-N,3-dimethyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-53-5 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]methylamino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]-N,3-dimethyl-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-55-7 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-56-8 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]methylamino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-58-0 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]methylamino]-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]-N,3-dimethyl-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-59-1 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-amino-3-(4-fluorophenyl)propyl]-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-60-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-61-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-62-6 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-63-7 HCAPLUS


CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-

Updated Search

09890219

dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-64-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-65-9 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287206-66-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,α-diethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-67-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,α-diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-68-2 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,α-diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-69-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-70-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-71-7 HCAPLUS


CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-

Updated Search

09890219

dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-72-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-73-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287206-74-0 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-75-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-76-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-77-3 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,Nα-diethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-78-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-propyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-79-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(1-methylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-80-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-N-cyclopropyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-81-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-82-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-83-1 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-84-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-85-3 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-86-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-87-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-88-6 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-89-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-90-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-91-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287206-92-2 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-(4-morpholinyl)-2-oxoethyl]-N,N2-dimethyl- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 287206-93-3 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[4-(methylsulfonyl)-1-piperazinyl]-2-oxoethyl]-N,N2-dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287206-94-4 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[4-(2-ethoxy-2-oxoethyl)-1-piperazinyl]-2-oxoethyl]-N,N2-dimethyl- (9CI) (CA INDEX NAME)

Updated Search

09890219


Absolute stereochemistry.



RN 287206-95-5 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1S)-2-[4-(carboxymethyl)-1-piperazinyl]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-oxoethyl]-N,N2-dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-96-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-Nα-propyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287206-97-7 HCAPLUS


CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2S)-2-(methylamino)butanoyl-3-

Updated Search

09890219

(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-98-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2R)-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-99-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-norvalyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287207-00-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-D-norvalyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-01-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-isoleucyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-02-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-D-isoleucyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287207-03-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-leucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-04-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-D-leucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287207-05-0 HCAPLUS


CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-4,5-didehydro-N-methyl-L-norvalyl-

Updated Search

09890219

3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-06-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-4,5-didehydro-N-methyl-D-norvalyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-07-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N,4-dimethyl-L-leucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287207-08-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N,4-dimethyl-D-leucyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-09-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2S)-4,4,4-trifluoro-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-10-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2S)-2-cyclohexyl-N-methylglycyl-3-(1,1-dimethylethyl)- α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


09890219



RN 287207-11-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2R)-2-cyclohexyl-N-methylglycyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-12-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-3-cyclohexyl-N-methyl-L-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-13-0 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-3-cyclohexyl-N-methyl-D-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-14-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-15-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-D-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287207-16-3 HCAPLUS


CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-4-fluoro-N-methyl-L-phenylalanyl-3-

Updated Search

09890219

(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-17-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-4-fluoro-N-methyl-D-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-18-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-4-chloro-N-methyl-L-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287207-19-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-4-chloro-N-methyl-D-phenylalanyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-20-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-tyrosyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287207-21-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-D-tyrosyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-22-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-3-(2-thienyl)-L-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-23-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-3-(2-thienyl)-D-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287207-24-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-3-cyclopropyl-N-methyl-L-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-25-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2S)-N-methyl-2-phenylglycyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287207-26-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N, α -dimethyl-L-phenylalanyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-27-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N, α -dimethyl-D-phenylalanyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-28-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N,2-dimethyl-L-leucyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-29-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-D-isovalyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287207-30-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N,3-dimethyl-D-isovalyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-31-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-1-(methylamino)cyclopentanecarbonyl-1-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-32-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-1-(methylamino)cyclohexanecarbonyl-1-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287207-33-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N,3-dimethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-34-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N,3-dimethyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287207-35-6 HCAPLUS


CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2R)-N-methyl-2-phenylglycyl-3-

Updated Search

09890219

(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-38-9 HCAPLUS

CN L-Tyrosinamide, 4-(trifluoromethyl)-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-39-0 HCAPLUS

CN L-Tyrosinamide, N-[(2R)-3-(4-fluorophenyl)-2-(hydroxymethyl)-1-oxopropyl]-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-41-4 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-cyano-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-31-5 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1R)-1-(1,4-dihydro-6-methyl-4-oxo-2-pyrimidinyl)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-73-5 HCAPLUS

CN Butanamide, 2-[[[(2S)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287209-74-9 HCAPLUS

CN Butanamide, 2-[[(2R)-2-amino-3-(4-fluorophenyl)propyl]amino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]-3-methyl-, (2S)- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 287209-77-2 HCAPLUS

CN Butanamide, 2-[[(2R)-2-amino-3-(4-fluorophenyl)propyl]methylamino]-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]-3-methyl-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287211-41-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-(2R)-4,4,4-trifluoro-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287212-51-5 HCAPLUS

Updated Search

09890219


CN L-Tyrosinamide, N-[(2S)-3-(4-fluorophenyl)-2-(hydroxymethyl)-1-oxopropyl]-
N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.




RN 287212-55-9 HCAPLUS

CN Tyrosinamide, 4-fluorophenylalanyl-N-methylvalyl-3-(1,1-dimethylethyl)-N-2-pyridinyl- (9CI) (CA INDEX NAME)



RN 287212-56-0 HCAPLUS

CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[(aminocarbonyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)




RN 287212-57-1 HCAPLUS

CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[(aminosulfonyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA

Updated Search

09890219


INDEX NAME)



RN 287212-58-2 HCAPLUS


CN Valinamide, 4-fluoro-L-phenylalanyl-N-[1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[(methylsulfonyl)amino]ethyl]-N2-methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



RN 287212-59-3 HCAPLUS

CN Valinamide, N-(4-fluorophenyl)alanyl-N-[3-amino-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-3-oxopropyl]-N2-methyl- (9CI) (CA INDEX NAME)




RN 287212-60-6 HCAPLUS


CN Valinamide, N-(4-fluorophenyl)alanyl-N-[1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-(methylsulfonyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Updated Search


09890219




RN 287212-61-7 HCAPLUS
 CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)



RN 287212-62-8 HCAPLUS
 CN Valinamide, N-(4-fluorophenyl)alanyl-N-[1-(1,4-dihydro-6-methyl-4-oxo-2-pyrimidinyl)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)



RN 287212-63-9 HCAPLUS
 CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1,3,4-oxadiazol-2-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)




Updated Search

09890219


RN 287212-64-0 HCAPLUS

CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1,2,4-oxadiazol-5-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)




RN 287212-65-1 HCAPLUS

CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)



RN 287212-66-2 HCAPLUS

CN Valinamide, N-(4-fluorophenyl)alanyl-N-[2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1H-1,2,4-triazol-3-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)



Updated Search

09890219

IT 287207-47-0P 287207-48-1P 287207-49-2P
287207-50-5P 287207-51-6P 287207-55-0P
287207-60-7P 287207-66-3P 287207-72-1P
287207-78-7P 287207-81-2P 287207-82-3P
287207-84-5P 287207-86-7P 287207-88-9P
287207-90-3P 287207-98-1P 287208-07-5P
287208-13-3P 287208-16-6P 287208-20-2P
287208-29-1P 287208-30-4P 287208-35-9P
287208-42-8P 287208-45-1P 287208-47-3P
287208-61-1P 287208-63-3P 287208-65-5P
287208-67-7P 287208-71-3P 287208-72-4P
287208-74-6P 287208-75-7P 287208-93-9P
287208-94-0P 287208-97-3P 287208-98-4P
287208-99-5P 287209-00-1P 287209-01-2P
287209-04-5P 287209-05-6P 287209-06-7P
287209-09-0P 287209-10-3P 287209-11-4P
287209-14-7P 287209-15-8P 287209-16-9P
287209-19-2P 287209-20-5P 287209-21-6P
287209-24-9P 287209-25-0P 287209-26-1P
287209-29-4P 287209-30-7P 287209-31-8P
287209-34-1P 287209-35-2P 287209-36-3P
287209-39-6P 287209-40-9P 287209-41-0P
287209-44-3P 287209-45-4P 287209-46-5P
287209-49-8P 287209-52-3P 287209-54-5P
287209-55-6P 287209-60-3P 287209-63-6P
287209-64-7P 287209-67-0P 287209-68-1P
287209-72-7P 287209-73-8P 287209-75-0P
287209-76-1P 287209-82-9P 287209-85-2P
287209-86-3P 287210-09-7P 287210-10-0P
287210-11-1P 287210-14-4P 287210-15-5P
287210-16-6P 287210-19-9P 287210-20-2P
287210-21-3P 287210-24-6P 287210-25-7P
287210-26-8P 287210-29-1P 287210-30-4P
287210-31-5P 287210-34-8P 287210-35-9P
287210-36-0P 287210-39-3P 287210-42-8P
287210-45-1P 287210-46-2P 287210-47-3P
287210-48-4P 287210-49-5P 287210-50-8P
287210-51-9P 287210-52-0P 287210-53-1P
287210-54-2P 287210-55-3P 287210-56-4P
287210-57-5P 287210-58-6P 287210-59-7P
287210-60-0P 287210-61-1P 287210-62-2P
287210-63-3P 287210-64-4P 287210-67-7P
287210-70-2P 287210-73-5P 287210-76-8P
287210-79-1P 287211-02-3P 287211-05-6P
287211-08-9P 287211-11-4P 287211-14-7P
287211-17-0P 287211-21-6P 287211-24-9P
287211-27-2P 287211-30-7P 287211-33-0P
287211-36-3P 287211-40-9P 287211-42-1P
287211-45-4P 287211-48-7P 287211-51-2P
287211-54-5P 287211-57-8P 287211-60-3P
287211-63-6P 287211-72-7P 287211-77-2P
287211-80-7P 287211-83-0P 287211-86-3P
287211-89-6P 287211-92-1P 287211-95-4P
287212-00-4P 287212-01-5P 287212-04-8P
287212-07-1P 287212-10-6P 287212-13-9P
287212-16-2P 287212-19-5P 287212-22-0P
287212-25-3P 287212-31-1P 287212-49-1P
287212-50-4P 287212-53-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT

Updated Search

09890219

(Reactant or reagent)


(preparation of peptides or analogs containing substituted phenethylamine moiety

as motilin receptor antagonists and drugs for preventing digestive tract movement or high level of blood motilin)

RN 287207-47-0 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-48-1 HCAPLUS

CN L-Tyrosinamide, 4-chloro-N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287207-49-2 HCAPLUS


CN L-Tyrosinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]-3,4-difluoro-L-

Updated Search

09890219

phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 287207-50-5 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-3-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287207-51-6 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-2-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287207-55-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl-N-(methanesulfonyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-60-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-methoxy-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

RN 287207-66-3 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy) carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-2-pyridinyl- (9CI) (CA INDEX NAME)

CC(C)S(=O)(=O)N[C@@H](Cc1ccc(O)c(Cc2ccc(O)c(C(=O)Nc3ccncc3)c2)c1)C(=O)N[C@@H](Cc1ccc(F)cc1)C(=O)NOC(C)(C)C


Absolute stereochemistry.



Absolute stereochemistry.

Updated Search


09890219



RN 287207-81-2 HCAPLUS

CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-2-amino-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-82-3 HCAPLUS

CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-2-[(aminoiminomethyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287207-84-5 HCAPLUS

CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-2-[[[(cyanoamino)(methylamino)methylene]amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-86-7 HCAPLUS

CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-2-[(aminosulfonyl)amino]-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287207-88-9 HCAPLUS

CN Glycine, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-L-tyrosyl-ψ(CH2-NH)-, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287207-90-3 HCAPLUS

20/207 90 9 NCI 100
CN Glycinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-
L-valyl-3-(1,1-dimethylethyl)-L-tyrosyl-ψ(CH₂-NH)- (9CI) (CA INDEX
NAME)


Absolute stereochemistry.



RN 287207-98-1 HCAPLUS

28/207 90 1 NCM E05
CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[(methylsulfonyl)amino]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-07-5 HCAPLUS

L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-3-amino-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-3-oxopropyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287208-13-3 HCAPLUS

CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-(methylsulfonyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-16-6 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-20-2 HCAPLUS

CN 2-Thia-5,8,11-triazadodecan-12-oic acid, 4-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-10-[(4-fluorophenyl)methyl]-7-(1-methylethyl)-6-oxo-, 1,1-dimethylethyl ester, 2,2-dioxide, (4S,7S,10S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219


RN 287208-29-1 HCAPLUS
CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-
[(1S)-1-(1,4-dihydro-6-methyl-4-oxo-2-pyrimidinyl)-2-[3-(1,1-
dimethylethyl)-4-hydroxyphenyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287208-30-4 HCAPLUS
CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-
[(1R)-1-(1,4-dihydro-6-methyl-4-oxo-2-pyrimidinyl)-2-[3-(1,1-
dimethylethyl)-4-hydroxyphenyl]ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287208-35-9 HCAPLUS
CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-
2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2,5-dioxo-4-
imidazolidinyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287208-42-8 HCAPLUS
CN L-Valinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1,3,4-oxadiazol-2-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287208-45-1 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287208-47-3 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1,2,4-oxadiazol-5-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-61-1 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287208-63-3 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-65-5 HCAPLUS

CN L-Valinamide, 4-fluoro-L-phenylalanyl-N-[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287208-67-7 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(1H-1,2,4-triazol-3-yl)ethyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287208-71-3 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]amino]carbonyl]-2-methylpropyl]amino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-72-4 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(2-thiazolyl)ethyl]amino]carbonyl]-2-methylpropyl]amino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287208-74-6 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-2-fluoro-L-tyrosyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287208-75-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-3-fluoro-L-tyrosyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287208-93-9 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287208-94-0 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287208-97-3 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

CC(C)C1=CC=C(C=C1)C(=O)N[C@@H](C)C(=O)S[C@@H](C(=O)N[C@@H](Cc1ccc(O)cc1)C(=O)N)C(=O)S[C@@H](Cc1ccc(F)cc1)C(=O)N

Absolute stereochemistry.



Absolute stereochemistry.




Updated Search

09890219

RN 287209-00-1 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 287209-01-2 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287209-04-5 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219




RN 287209-05-6 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287209-06-7 HCAPLUS
CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -dimethyl-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287209-09-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-10-3 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-11-4 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-14-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-15-8 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287209-16-9 HCAPLUS
CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-19-2 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-20-5 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-21-6 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-24-9 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-25-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219




RN 287209-26-1 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-N-ethyl-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287209-29-4 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-30-7 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-31-8 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287209-34-1 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N-dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-35-2 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,Na-dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-36-3 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,Na-dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287209-39-6 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-40-9 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-41-0 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-44-3 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-45-4 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-46-5 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-49-8 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-N,3-bis(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-52-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl-N-[(methylsulfonyl)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-54-5 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-(4-fluorophenyl)propyl]-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-55-6 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-(4-fluorophenyl)propyl]-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-60-3 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-(4-fluorophenyl)propyl]-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287209-63-6 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]amino]carbonyl]-2-methylpropyl]amino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-64-7 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]amino]carbonyl]-2-methylpropyl]methylamino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-67-0 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]methylamino]carbonyl]-2-methylpropyl]amino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287209-68-1 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]methylamino]carbonyl]-2-methylpropyl]methylamino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-72-7 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]amino]carbonyl]-2-methylpropyl]amino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-73-8 HCAPLUS

CN Carbamic acid, [(1R)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]amino]carbonyl]-2-methylpropyl]amino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287209-75-0 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]amino]carbonyl]-2-methylpropyl]methylamino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-76-1 HCAPLUS

CN Carbamic acid, [(1R)-2-[[[(1S)-1-[[[(1R)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-methylethyl]amino]carbonyl]-2-methylpropyl]methylamino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287209-82-9 HCAPLUS

CN Carbamic acid, [(1S)-2-[[[(1S)-1-[[[(1S)-2-[3-(1,1-dimethylethyl)-4-hydroxyphenyl]-1-(hydroxymethyl)ethyl]methylamino]carbonyl]-2-methylpropyl]methylamino]-1-[(4-fluorophenyl)methyl]ethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287209-85-2 HCAPLUS

L-Tyrosinamide, N-[(2S)-2-[[(1,1-dimethylethoxy) carbonyl] amino]-3-(4-fluorophenyl)propyl]-L-valyl-N-[(1,1-dimethylethoxy) carbonyl]-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287209-86-3 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-2-[[[(1,1-dimethylethoxy)carbonyl]amino]-3-(4-fluorophenyl)propyl]-N-methyl-L-valyl-N-[(1,1-dimethylethoxy)carbonyl]-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-09-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219




RN 287210-10-0 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287210-11-1 HCAPLUS
CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287210-14-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-15-5 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-16-6 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-19-9 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy) carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-20-2 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy) carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN	287210-21-3	HCAPLUS
CN	L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-N-ethyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N,Nα-diethyl-(9CI) (CA INDEX NAME)	

Absolute stereochemistry.



```


RN      287210-24-6  HCAPLUS
CN      L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-
        ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI)  (CA INDEX NAME)

```

Absolute stereochemistry.

Updated Search


09890219



RN 287210-25-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-26-8 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287210-29-1 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-30-4 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-31-5 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-34-8 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-35-9 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-36-0 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N,N α -diethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-39-3 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-propyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-42-8 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(1-methylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-45-1 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-N-cyclopropyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-46-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-47-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287210-48-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search

Chemical structure of compound 10: A bis-thioamide derivative. It features two 4-fluorophenyl groups and one 3-(tert-butyl)-4-hydroxyphenyl group. The central core consists of two thioamide groups linked by a carbonyl group. The left thioamide is substituted with a 4-fluorophenyl group and a 1-methyl-2-((benzyloxy)carbonyl)aminoethyl group. The right thioamide is substituted with a 3-(tert-butyl)-4-hydroxyphenyl group and an isopropyl group. A benzyl group is attached to the carbonyl oxygen of the left thioamide.

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-(9CI) (CA INDEX NAME)

Chemical structure of compound 10, a cyclic peptide derivative. The structure shows a central six-membered ring containing two sulfur atoms and two nitrogen atoms. The ring is substituted with a 4-fluorophenyl group, a 4-hydroxy-3-tert-butylphenyl group, and a 2-methyl-2-isopropyl-1-oxoethyl group. A side chain includes a benzyl group and a hydroxymethyl group.

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Chemical structure of compound 10, a 1,3,5-trithiane derivative. The structure shows a central 1,3,5-trithiane ring with a methyl group on the nitrogen. The sulfur atoms are substituted with a 4-(tert-butyl)phenyl group, a 4-fluorophenyl group, and a 2-((S)-1-aminopropyl)acetyl group. The amino group is shown as H₂N.


Updated Search

09890219

RN 287210-51-9 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-52-0 HCAPLUS

L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-53-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-54-2 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 287210-55-3 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-56-4 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-57-5 HCAPLUS

207210 37 3 NCM 208
CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-
methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-58-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287210-59-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-Nα-ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-60-0 HCAPLUS

CN L-Tyrosinamide, N-ethyl-4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-Nα-ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-61-1 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-62-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-63-3 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-64-4 HCAPLUS

Updated Search

09890219

CN L-Tyrosinamide, 4-fluoro-N-methyl-N-[(phenylmethoxy) carbonyl]-L-phenylalanyl-N-ethyl-L-valyl-3-(1,1-dimethylethyl)-N α -ethyl-N-(hydroxymethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-67-7 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy) carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-(hydroxymethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-70-2 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy) carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-(4-morpholinyl)-2-oxoethyl]-N,N2-dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287210-73-5 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[4-(methylsulfonyl)-1-piperazinyl]-2-oxoethyl]-N,N2-dimethyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287210-76-8 HCAPLUS

CN L-Valinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-[(1S)-1-[[3-(1,1-dimethylethyl)-4-hydroxyphenyl]methyl]-2-[4-(2-ethoxy-2-oxoethyl)-1-piperazinyl]-2-oxoethyl]-N,N2-dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287210-79-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -(1-methylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287211-02-3 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-
(2S)-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287211-05-6 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-
(2R)-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287211-08-9 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-norvalyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-11-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-D-norvalyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287211-14-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-isoleucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287211-17-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-D-isoleucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287211-21-6 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-leucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287211-24-9 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-D-leucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-27-2 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-4,5-didehydro-N-methyl-L-norvalyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287211-30-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-4,5-didehydro-N-methyl-D-norvalyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-33-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N,4-dimethyl-L-leucyl-3-(1,1-dimethylethyl)-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287211-36-3 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-
N,4-dimethyl-D-leucyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA
INDEX NAME)

Absolute stereochemistry.




RN 287211-40-9 HCAPLUS
CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-(2S)-
4,4,4-trifluoro-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)-N α -
methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287211-42-1 HCAPLUS
CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-(2R)-
4,4,4-trifluoro-2-(methylamino)butanoyl-3-(1,1-dimethylethyl)-N α -
methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-45-4 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-
(2S)-2-cyclohexyl-N-methylglycyl-3-(1,1-dimethylethyl)-N α -methyl-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287211-48-7 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-(2R)-2-cyclohexyl-N-methylglycyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-51-2 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-3-cyclohexyl-N-methyl-L-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287211-54-5 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-3-cyclohexyl-N-methyl-D-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287211-57-8 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219




RN 287211-60-3 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-D-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 287211-63-6 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-4-fluoro-N-methyl-L-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287211-72-7 HCAPLUS

L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-4-fluoro-N-methyl-D-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl-
 (9CI): (CA INDEX NAME)


Absolute stereochemistry.



RN 287211-77-2 HCAPLUS

L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-4-chloro-N-methyl-L-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-80-7 HCAPLUS

287211 00 1 NON-188
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-4-chloro-N-methyl-D-phenylalanyl-3-(1,1-dimethylethyl)-N α -methyl-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287211-83-0 HCAPLUS
CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-tyrosyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287211-86-3 HCAPLUS
CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N-methyl-D-tyrosyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Updated Search

09890219



RN 287211-89-6 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-3-(2-thienyl)-L-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287211-92-1 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-3-(2-thienyl)-D-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287211-95-4 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-3-cyclopropyl-N-methyl-L-alanyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287212-00-4 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-(2S)-N-methyl-2-phenylglycyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287212-01-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-(2R)-N-methyl-2-phenylglycyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287212-04-8 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-
N, α -dimethyl-L-phenylalanyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.




RN 287212-07-1 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-
N, α -dimethyl-D-phenylalanyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX
NAME)

Updated Search

Absolute stereochemistry.




RN	287212-13-9	HCAPLUS
CN	L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-D-isovalyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)	

RN	287212-16-2	HCAPLUS
CN	L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N,3-dimethyl-D-isovalyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)	

Updated Search


09890219



RN 287212-19-5 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-1-(methylamino)cyclopentanecarbonyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287212-22-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-1-(methylamino)cyclohexanecarbonyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




Updated Search

09890219

RN 287212-25-3 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N,3-dimethyl-L-valyl-3-(1,1-dimethylethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287212-31-1 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-N,3-dimethyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl-O-(phenylmethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 287212-49-1 HCAPLUS

CN L-Tyrosinamide, N-[(2S)-3-ethoxy-2-[(4-fluorophenyl)methyl]-1,3-dioxopropyl]-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search


09890219



RN 287212-50-4 HCAPLUS

CN L-Tyrosinamide, N-[(2R)-3-ethoxy-2-[(4-fluorophenyl)methyl]-1,3-dioxopropyl]-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.




RN 287212-53-7 HCAPLUS

CN L-Tyrosinamide, 4-cyano-N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

09890219



REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his

(FILE 'HOME' ENTERED AT 12:21:47 ON 11 JUN 2007)

FILE 'REGISTRY' ENTERED AT 12:21:55 ON 11 JUN 2007

L1 STRUCTURE UPLOADED

L2 1 S L1

L3 519 S L1 FULL

FILE 'HCAPLUS' ENTERED AT 12:27:12 ON 11 JUN 2007

L4 94 S L3

L5 2 S L4 AND MATSUOKA, H?/AU

=> s 14 not 15

L6 92 L4 NOT L5

=> s 16 and sato, t?/au

24044 SATO, T?/AU

L7 1 L6 AND SATO, T?/AU

=> d 17, ibib abs hitstr, 1

L7 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:90066 HCAPLUS

DOCUMENT NUMBER: 136:135034

TITLE: Method for producing tripeptide derivative

INVENTOR(S): Sato, Tsutomu; Shimizu, Hirohito

PATENT ASSIGNEE(S): Chugai Seiyaku Kabushiki Kaisha, Japan

SOURCE: PCT Int. Appl., 50 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.

KIND

DATE

APPLICATION NO.

DATE

Updated Search

```

-----
WO 2002008248      A1      20020131      WO 2001-JP6295      20010719
W:  AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
    CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
    GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
    LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,
    RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
    UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW:  GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
    DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
    BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
JP 2005097119      A      20050414      JP 2000-219977      20000721
PRIORITY APPLN. INFO.:      JP 2000-219977      A      20000721
OTHER SOURCE(S):      CASREACT 136:135034; MARPAT 136:135034
GI

```

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A method for producing L-phenylalanyl-L-valyl-L-3-tert-butyl-L-tyrosinamide compds. represented by the general formula (I; wherein R1 represents a hydrogen atom or a linear or branched aliphatic alkyl group having 1 to 4 carbon atoms; R2 represents a hydrogen atom or Me group; R3 represents a hydrogen atom or Me group; and R4 represents a halogen atom) comprises condensation of 3-tert-butyl-L-tyrosinamide derivs. (II; R1, R2 = same as above) with N-methyl-L-valine derivs. (III; P1 = amino-protecting group), N-deprotection of the resulting L-valyl-3-tert-butyl-L-tyrosinamide derivs. (IV; R1, R2, P1 = same as above), and condensation of the resulting IV (P1 = H; R1, R2 = same as above) with L-phenylalanine derivs. (V; R3, R4 = same as above; P2 = amino-protecting group) followed by N-deprotection. The method can be advantageously used for producing a novel peptide derivative in a com. process. Thus, 20.8 g MeSO3H and 20.0 g tert-Bu chloride were successively added to 10.0 g L-tyrosine Me ester hydrochloride under stirring, stirred at 50° for 5 h, treated dropwise with MeOH (20 mL)/H2O (20 mL) under ice-cooling then with a solution of 14.2 g KOH in 43 mL H2O at <10° to give 77.0% 3-tert-butyl-L-tyrosine Me ester which (8.35 g) was added to a mixture of 24.1 g 62% aqueous ethylamine and 7.52 g ethylamine hydrochloride under ice-cooling and stirred at room temperature for


5 h to give 89.8% 3-tert-butyl-L-tyrosine ethylamide (VI). To a solution of 5.50 g VI and 3.35 g 1-hydroxybenzotriazole monohydrate in 55 mL THF were successively added 4.19 g 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and 3.04 mL Et3N and stirred at room temperature for 2.5 h to

give 100% N-tert-butoxycarbonyl-N-methyl-L-valyl-3-tert-butyl-L-tyrosine ethylamide which (10.0 g) was dissolved in 100 mL EtOAc, treated with 11.1 mL concentrated H2SO4 under ice-cooling, treated with 100 mL EtOAc, adjusted pH 8 by adding saturated aqueous NaHCO3, and stirred 15 min to give 87.9% N-methyl-L-valyl-3-tert-butyl-L-tyrosine ethylamide (VII). To a mixture of 5.50 g VII, 5.20 g N-tert-butoxycarbonyl-N-methyl-4-fluoro-L-phenylalanine, 4.47 g 2-chloro-1-methylpyridinium iodide, and 37 mL tert-Bu Me ether was added 5.09 mL Et3N and stirred at room temperature for 4 h to give 86.0% N-tert-butoxycarbonyl-N-methyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-tert-butyl-L-tyrosine ethylamide which (7.50 g) was similarly deprotected as described above using concentrated H2SO4 in EtOAc to give 100% N-methyl-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-tert-butyl-L-

09890219


tyrosine.
IT 287210-10-0P 393562-03-3P
RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (preparation tripeptide derivs. by sequential coupling of N-methyl-L-valine derivs. and L-phenylalanine derivs. to 3-tert-butyl-L-tyrosinamide derivs.)
RN 287210-10-0 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 393562-03-3 HCAPLUS
CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N- α -methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



Updated Search

09890219

IT 287205-81-6P 287206-61-5P


RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)

(preparation tripeptide derivs. by sequential coupling of N-methyl-L-valine derivs. and L-phenylalanine derivs. to 3-tert-butyl-L-tyrosinamide derivs.)

RN 287205-81-6 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N α -methyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 287206-61-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

7

THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his

(FILE 'HOME' ENTERED AT 12:21:47 ON 11 JUN 2007)

FILE 'REGISTRY' ENTERED AT 12:21:55 ON 11 JUN 2007

L1 STRUCTURE UPLOADED

L2 1 S L1

Updated Search

09890219

L3 519 S L1 FULL

FILE 'HCAPLUS' ENTERED AT 12:27:12 ON 11 JUN 2007

L4 94 S L3
L5 2 S L4 AND MATSUOKA, H?/AU
L6 92 S L4 NOT L5
L7 1 S L6 AND SATO, T?/AU

=> s 16 not 17

L8 91 L6 NOT L7

=> s 18 and takahashi, t?/au

20393 TAKAHASHI, T?/AU

L9 0 L8 AND TAKAHASHI, T?/AU

=> s 18 and kim, d?/au

24688 KIM, D?/AU

L10 1 L8 AND KIM, D?/AU

=> d 110, ibib abs hitstr, 1

L10 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:637704 HCAPLUS

DOCUMENT NUMBER: 137:185838

TITLE: Process for preparation of peptide derivatives

INVENTOR(S): Kim, Dong Ick; Jeon, Gee Ho; Kim, Sung Jin

PATENT ASSIGNEE(S): Chugai Seiyaku Kabushiki Kaisha, Japan

SOURCE: PCT Int. Appl., 40 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002064623	A1	20020822	WO 2002-JP1139	20020212
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2002230216	A1	20020828	AU 2002-230216	20020212
PRIORITY APPLN. INFO.:			KR 2001-6673	A 20010212
			WO 2002-JP1139	W 20020212
OTHER SOURCE(S):			CASREACT 137:185838; MARPAT 137:185838	
GI				



AB The title compds. I [R1 is hydrogen or linear or branched C1-4 alkyl; R2 is hydrogen or linear or branched C1-4 alkyl; and R3 is halogeno] are prepared in a multistep process. I are motilin receptor antagonists and are useful as drugs for gastric or intestinal diseases (no data). Thus, amidation of N-(tert-butoxycarbonyl)-L-(4-fluorophenyl)alanine with L-valine Me ester hydrochloride, followed by methylation with iodomethane, saponification, reaction with 3-tert-butyl-L-tyrosine Et amide, and deprotection,

gave N-methyl-L-4-fluorophenylalanyl-N-methyl-L-valine-3-tert-butyl-L-tyrosine Et amide.

IT 287206-61-5P


RL: IMF (Industrial manufacture); PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(process for preparation of peptide derivs.)

RN 287206-61-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 287210-10-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(process for preparation of peptide derivs.)

RN 287210-10-0 HCAPLUS

CN L-Tyrosinamide, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl- (9CI) (CA INDEX NAME)

09890219

Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his

{FILE 'HOME' ENTERED AT 12:21:47 ON 11 JUN 2007}

FILE 'REGISTRY' ENTERED AT 12:21:55 ON 11 JUN 2007

L1 STRUCTURE UPLOADED
L2 1 S L1
L3 519 S L1 FULL

FILE 'HCAPLUS' ENTERED AT 12:27:12 ON 11 JUN 2007

L4 94 S L3
L5 2 S L4 AND MATSUOKA, H?/AU
L6 92 S L4 NOT L5
L7 1 S L6 AND SATO, T?/AU
L8 91 S L6 NOT L7
L9 0 S L8 AND TAKAHASHI, T?/AU
L10 1 S L8 AND KIM, D?/AU

=> s l8 not l10

L11 90 L8 NOT L10

=> s l11 and jung, k?/au

2843 JUNG, K?/AU

L12 0 L11 AND JUNG, K?/AU

=> s l11 and park, c?/au

10333 PARK, C?/AU

L13 0 L11 AND PARK, C?/AU

=> d l11, ibib abs fhitr, 1-90

L11 ANSWER 1 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2007:16167 HCAPLUS

DOCUMENT NUMBER: 146:252099

Updated Search

09890219

TITLE: Oxidative cyclization of isodityrosine tripeptides:
optimized condition and application of
electrochemically generated thallium(III) ion

AUTHOR(S): Tanabe, Takamasa; Obata, Rika; Nishiyama, Shigeru

CORPORATE SOURCE: Department of Chemistry, Faculty of Science and
Technology, Keio University, Hiyoshi 3-14-1,
Kohoku-ku, Yokohama, 223-8522, Japan

SOURCE: Heterocycles (2006), 69, 113-118
CODEN: HTCYAM; ISSN: 0385-5414

PUBLISHER: Japan Institute of Heterocyclic Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 146:252099


AB Thallium(III) trinitrate oxidation of the tripeptide, which was a synthetic
precursor of the 17-membered isodityrosine natural products isolated from
sponge Microcionia eurya, euryamides, was investigated to give the
desired cyclized compound in 96% yield at best. In addition, the thallium(III)
species generated by electrochem. oxidation of thallium(I) successfully
produced the target compds.

IT 866417-84-7
RL: RCT (Reactant); RACT (Reactant or reagent)
(oxidative macrocyclization of isodityrosine tripeptides by
electrochem. generated thallium)

RN 866417-84-7 HCAPLUS

CN L-Tyrosine, 3,5-diiodo-N-[(phenylmethoxy)carbonyl]-L-tyrosyl-O-
(phenylmethyl)-L-threonyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX
NAME)

Absolute stereochemistry.



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 2 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1298385 HCAPLUS

DOCUMENT NUMBER: 146:177451

TITLE: Delineation of the motilin domain involved in
desensitization and internalization of the motilin
receptor by using full and partial antagonists

AUTHOR(S): Mitselos, Anna; Depoortere, Inge; Peeters, Theo L.

CORPORATE SOURCE: Centre for Gastroenterological Research, Catholic
University of Leuven, Louvain, B-3000, Belg.

SOURCE: Biochemical Pharmacology (2007), 73(1), 115-124

CODEN: BCPA6; ISSN: 0006-2952

PUBLISHER: Elsevier B.V.

Updated Search

DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Studies with fragments of the gastrointestinal peptide, motilin, indicate that the C-terminal region of this peptide plays an important role in the desensitization of the motilin receptor (MTLR). To verify this hypothesis, we studied the desensitization, phosphorylation and internalization induced by motilin analogs of different chain length with agonistic and antagonistic properties in CHO-MTLR cells. We studied motilin [1-22], the [1-14] fragment, the analogs Phe3[1-22] and Phe3[1-14], and two putative antagonists, GM-109 and MA-2029 (modified 1-4 and 1-3 fragments). Activation and desensitization (2 h preincubation with the motilin analogs 10 μ M) were studied in CHO-MTLR cells by an aequorin based luminescence assay. Phosphorylation was studied by immunopptn. and internalization was visualized in CHO-MTLR cells containing an enhanced green fluorescent protein (CHO-MTLR-EGFP). Results showed that Motilin [1-22] and [1-14] were more potent than Phe3[1-22] and Phe3[1-14] (pEC50: 9.77, 8.78, 7.36 and 6.65, resp.) to induce Ca²⁺ release. GM-109 and MA-2029 were without agonist activity. Motilin[1-22] and Phe3[1-22] decreased the second response to motilin from 78 \pm 2% to 11 \pm 3% and 34 \pm 3% (P < 0.001), resp., whereas [1-14], Phe3[1-14], GM-109 and MA-2029 had no desensitizing effect (68 \pm 5%, 78 \pm 3%, 78 \pm 6% and 78 \pm 5%, resp., P > 0.05). The rank order of MTLR-phosphorylation was: [1-22] > [1-14] > Phe3[1-22] = Phe3[1-14] > GM-109 = MA-2029. Only motilin [1-22] and [1-14] induced receptor MTLR-EGFP internalization as shown by a decrease in membrane fluorescence: 20 \pm 3% and 7 \pm 3%, resp. Thus, the C-terminus of motilin enhances desensitization, phosphorylation and internalization of the MTLR while modifications of the N-terminus can favor a conformation of the receptor that is less susceptible to phosphorylation and internalization.

IT 922190-03-2, MA 2029


RL: BSU (Biological study, unclassified); PKT (Pharmacokinetics); PRP (Properties); BIOL (Biological study)

(motilin receptor antagonist; delineation of motilin domain involved in desensitization, phosphorylation and internalization of motilin receptor by using full and partial antagonists)

RN 922190-03-2 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-methyl-L-phenylalanyl-N-methyl-L-valyl-3-(1,1-dimethylethyl)-N-ethyl-, hydrochloride (1:1) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

09890219

REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:710144 HCAPLUS

DOCUMENT NUMBER: 145:189182

TITLE: Preparation of peptides as neuropeptide-2 receptor (Y-2R) agonists

INVENTOR(S): Danho, Waleed; Ehrlich, George; Fry, David C.; Khan, Wajiha; Swistok, Joseph

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 102 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2006160742	A1	20060720	US 2006-328743	20060110
WO 2006077035	A1	20060727	WO 2006-EP161	20060111
WO 2006077035	A9	20061026		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			

PRIORITY APPLN. INFO.: US 2005-644840P P 20050118

OTHER SOURCE(S): MARPAT 145:189182

AB The invention relates to peptides Y-R1-R2-X-R3-R4-R5-R6-R7-R8-R9-R10-R11-R12-R13-R14-NH2 [X is 6-piperazin-1-yl-4(3H)-quinazolinone-3-acetic acid (Pqa), 5-O-(carboxymethyl)serotonin (Cms), 4-(2-aminoethyl)-6-dibenzofuranpropanoic acid, 4-piperidin-4-ylbutanoic acid, or 4-(2-aminoethyl)-1-(carboxymethyl)piperazine; Y is H, (un)substituted alkyl, aryl, or alkoxy or a poly(ethylene) glycol moiety; R1-R14 are amino acid residues (defined)] and their pharmaceutically-acceptable salts which are neuropeptide-2 receptor (Y-2R) agonists and are useful for the treatment of diseases such as obesity. Thus, H-Ile-Lys-Pqa-Arg-His-Tyr-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH2 was prepared by the solid-phase method, exhibited selective Y-2R activity in vitro (EC50 = 3.2 nM, IC50 = 0.032 nM), and was shown to cause reduction of food intake in mouse models of human obesity.

IT 900808-58-4P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of peptides as neuropeptide-2 receptor (Y-2R) agonists)

RN 900808-58-4 HCAPLUS

CN L-Tyrosinamide, N-acetyl-L-isoleucyl-L-norleucyl-4-oxo-6-(1-piperazinyl)-3(4H)-quinazolineacetyl-3,4,5-trifluoro-L-phenylalanyl-L-histidyl-L-tyrosyl-L-leucyl-L-asparaginyl-L-leucyl-L-valyl-L-threonyl-L-arginyl-L-


Updated Search

09890219


glutaminy-L-arginyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





L11 ANSWER 4 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:595922 HCAPLUS

DOCUMENT NUMBER: 146:179253

TITLE: TMC-95-Based Inhibitor Design Provides Evidence for the Catalytic Versatility of the Proteasome

AUTHOR(S): Groll, Michael; Goetz, Marion; Kaiser, Markus; Weyher, Elisabeth; Moroder, Luis

CORPORATE SOURCE: Department for Physiological Chemistry, Ludwig-Maximilians-University, Munich, D-81377, Germany

SOURCE: Chemistry & Biology (Cambridge, MA, United States) (2006), 13(6), 607-614

CODEN: CBOLE2; ISSN: 1074-5521

PUBLISHER: Cell Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB TMC-95's natural cyclic tripeptide metabolites represent potent competitive proteasome inhibitors. The constrained conformation of TMC-95 proteasomal inhibitors provides the driving force for entropically high-affinity binding. Based on the crystal structure of the proteasome:TMC-95A complex, the synthetically challenging TMC-95 core structure was used for the design and synthesis of less demanding biphenyl-ether macrocycles, in which the biphenyl-ether moiety functions as an endocyclic clamp restricting its tripeptide backbone. These simplified analogs allowed us to identify high plasticity of the proteasomal tryptic-like specificity pocket. Biphenyl-ether compds. extended with an amide group were hydrolyzed by the proteasome, although the crystal structure of such proteasome:biphenyl-ether complexes revealed quenching of proteolysis at the acyl-enzyme intermediate. Our data reveal that biphenyl-ether derivs. bind noncovalently to the proteasomal tryptic-like active site in a reversible substrate-like manner without allosteric changes of active site residues.

IT 918906-65-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(design, preparation and structural characterization of proteasomal TMC-95-based inhibitors)

RN 918906-65-7 HCAPLUS

CN L-Tyrosinamide, 3-fluoro-4-nitro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-L-asparaginyl-N-propyl- (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 5 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:505970 HCAPLUS

DOCUMENT NUMBER: 146:274598

TITLE: Solid-phase synthesis and biological activities of biphenyl ether-containing cyclic oligopeptides

AUTHOR(S): Nakamura, Kazuhiko; Obata, Rika; Nishiyama, Shigeru

CORPORATE SOURCE: Faculty of Science and Technology, Keio University, 3-14-1 Kohoku-ku, Yokohama, 223-8522, Japan

SOURCE: Peptide Science (2006), Volume Date 2005, 42nd, 127-130

CODEN: PSCIFQ; ISSN: 1344-7661

PUBLISHER: Japanese Peptide Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A symposium report. Halogenated cyclic isodityrosine-class tripeptides were synthesized as analogs of eurylamide B, a marine natural product. The cyclic peptides were synthesized using thallium(III) oxidation as a key reaction to construct the biaryl ether linkage in good yield. In addition, manipulation of trityl resin-supported tripeptide as a substrate of oxidation enabled a solid-phase synthesis of the peptides. Upon combination with imipenem, some of the resultant eurylamide-analogs showed anti-MRSA activity.

IT 866417-84-7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(solid-phase synthesis of biphenyl ether-linked cyclic peptides as eurylamide B analogs, and their anti-MRSA activity in combination with imipenem)

RN 866417-84-7 HCAPLUS

CN L-Tyrosine, 3,5-diiodo-N-[(phenylmethoxy)carbonyl]-L-tyrosyl-O-(phenylmethyl)-L-threonyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 6 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:494455 HCAPLUS

DOCUMENT NUMBER: 144:488943

TITLE: Preparation of cyclic isodityrosine derivatives having an imipenem-potentiating effect and a cholesteryl ester formation inhibiting activity

INVENTOR(S): Nishiyama, Shigeru; Obata, Rika; Tomoda, Hiroshi

PATENT ASSIGNEE(S): Keio University, Japan

SOURCE: PCT Int. Appl., 75 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006054396	A1	20060526	WO 2005-JP18182	20050930
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRIORITY APPLN. INFO.:

JP 2004-336698

A 20041119

OTHER SOURCE(S):

MARPAT 144:488943

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I [R1 = H, amido, Boc, etc.; R2 = benzoyl, benzyl, modified benzyl, etc.; R3 = H, OR5, amido; R5 = straight-chain or branched alkyl, aromatic ring; R4 = H, straight-chain or branched alkyl; X1, X2 = halo] and

their pharmacol. acceptable salts were prepared For example, BOP mediated acylation of 3,5-dibromo-L-tyrosine Me ester with O-benzyl-N-Boc-threonine followed by reaction with N-Cbz-3,5-diiodo-L-tyrosine and treatment with thallium (III) nitrate afforded compound II. Compound II exhibited an imipenem-potentiating effect against MRSA (IC₅₀ = 3 µg/mL). And, in cholesteryl ester formation inhibition assays, the IC₅₀ value of compound II was 3 µg/mL.

IT 866417-84-7P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of cyclic isodityrosine derivs. having imipenem-potentiating effect and cholesteryl ester formation inhibiting activity)

RN 866417-84-7 HCAPLUS

CN L-Tyrosine, 3,5-diiodo-N-[(phenylmethoxy)carbonyl]-L-tyrosyl-O-[(phenylmethyl)-L-threonyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 7 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:1007538 HCAPLUS

DOCUMENT NUMBER: 143:460423

TITLE: Further analogues of plant peptide hormone phytosulfokine- α (PSK- α) and their biological evaluation

AUTHOR(S): Bahyrycz, Agata; Matsubayashi, Yoshikatsu; Ogawa, Mari; Sakagami, Youji; Konopinska, Danuta

CORPORATE SOURCE: Faculty of Chemistry, University of Wroclaw, Wroclaw, 50-383, Pol.

SOURCE: Journal of Peptide Science (2005), 11(9), 589-592
CODEN: JPSIEI; ISSN: 1075-2617

PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:460423

AB Phytosulfokine- α (PSK- α), a sulfated growth factor of structure H-Tyr(SO₃H)-Ile-Tyr(SO₃H)-Thr-Gln-OH universally found in both monocotyledons and dicotyledons, strongly promotes proliferation of plant cells in culture. In studies on the structure/activity relationship of PSK- α , synthesis was performed for a series of 23 analogs modified at position 1, 3 or 4 as well as simultaneously at positions 1 and 3 of the peptide chain. The peptides were synthesized by the solid phase

09890219

method according to the Fmoc procedure on a Wang-resin. Free peptides were released from the resin by 95% TFA in the presence of EDT. All peptides were tested by competitive binding assay to the carrot membrane using ^3H -labeled PSK- α according to the test of Matsubayashi et al. Among these peptide analogs, [H-Phe(4-Cl)1]-PSK- α , [H-Phe(4-I)1]-PSK- α , and [Phe(4-Cl)3]-PSK- α retained 30% PSK- α activity. Analog [Tyr(PO $^3\text{H}_2$)3]-PSK- α showed 10% of PSK- α activity.


IT 869100-74-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation of analogs of plant peptide hormone phytosulfokine- α (PSK- α) and their biol. evaluation)

RN 869100-74-3 HCAPLUS

CN L-Glutamine, 4-chloro-L-phenylalanyl-L-isoleucyl-O-sulfo-L-tyrosyl-L-threonyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 8 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:921405 HCAPLUS

DOCUMENT NUMBER: 143:367587

TITLE: Discovery of halogenated euryamide B analogs as inhibitors of lipid droplet accumulation in macrophages

AUTHOR(S): Obata, Rika; Ohshiro, Taichi; Tomoda, Hiroshi; Nishiyama, Shigeru

CORPORATE SOURCE: Department of Chemistry, Faculty of Science and Technology, Keio University, Kohoku-ku, Yokohama, 223-8522, Japan


SOURCE: Bioorganic & Medicinal Chemistry Letters (2005), 15(19), 4189-4191

CODEN: BMCLE8; ISSN: 0960-894X

Updated Search

09890219

PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 143:367587
GI



AB Halogenated cyclic isodityrosine-tripeptides [I (R1 = H, R2 = CH2Ph, R3 = Me; R1 = R2 = H, R3 = Me; R1 = Boc, R2 = R3 = H; Boc = tert-butoxycarbonyl)] were synthesized as analogs of a marine natural product, eurypamide B. Although the original eurypamides showed no inhibitory activity, the new analogs were found to inhibit lipid droplet accumulation in macrophages with a low micromolar IC50 value.

IT 620960-60-3P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of halogenated cyclic isodityrosine-tripeptides eurypamide B analogs by peptide coupling and ring closure via intramol. phenolic oxidation)

RN 620960-60-3 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-3,5-diiodo-L-tyrosyl-L-threonyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT:

14

THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

Updated Search

09890219

L11 ANSWER 9 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:485754 HCAPLUS

DOCUMENT NUMBER: 143:173123

TITLE: Microwave-Assisted Intramolecular Suzuki-Miyaura Reaction to Macrocycle, a Concise Asymmetric Total Synthesis of Biphenomycin B

AUTHOR(S): Lepine, Renaud; Zhu, Jieping

CORPORATE SOURCE: Institut de Chimie des Substances Naturelles, CNRS, Gif-sur-Yvette, 91198, Fr.

SOURCE: Organic Letters (2005), 7(14), 2981-2984

CODEN: ORLEF7; ISSN: 1523-7060


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 143:173123

GI



AB A concise and efficient total synthesis of biphenomycin B (I) was accomplished featuring a key microwave-assisted intramol. Suzuki-Miyaura reaction for formation of the 15-membered meta,meta-cyclophane 20.


IT 861099-80-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(microwave-assisted intramol. Suzuki-Miyaura reaction to macrocycle, a concise asym. total synthesis of biphenomycin B)

RN 861099-80-1 HCAPLUS

CN L-Phenylalanine, N-[(1,1-dimethylethoxy)carbonyl]-5-iodo-2-(1-methylethoxy)-L-phenylalanyl-3-[(5R)-2,2-dimethyl-3-[(phenylmethoxy)carbonyl]-5-oxazolidinyl]-L-alanyl-2-(1-methylethoxy)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 66 THERE ARE 66 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 10 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:857446 HCAPLUS
 DOCUMENT NUMBER: 141:326194
 TITLE: Gonadotropin releasing hormone (GnRH) analogs
 conjugates with steroid hormones and therapeutic uses
 thereof
 INVENTOR(S): Millar, Robert Peter
 PATENT ASSIGNEE(S): Ardana Bioscience Limited, UK
 SOURCE: PCT Int. Appl., 76 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004087215	A1	20041014	WO 2004-GB1478	20040405
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1613357	A1	20060111	EP 2004-725716	20040405
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,			

09890219

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR
JP 2006522085 T 20060928 JP 2006-506091 20040405
US 2006247177 A1 20061102 US 2006-552110 20060703
PRIORITY APPLN. INFO.: GB 2003-7777 A 20030404
WO 2004-GB1478 W 20040405

AB A compound comprising a gonadotrophin releasing hormone analog conjugated to a hormone moiety, or a derivative thereof, which is able to bind to a plasma hormone binding protein. The compds. may be used to treat hormone-dependent disorders such as cancer, or as a contraceptive.

IT 428438-55-5, A-84861


RL: RCT (Reactant); RACT (Reactant or reagent)
(gonadotropin releasing hormone (GnRH) analogs conjugates with steroid hormones and therapeutic uses thereof)


RN 428438-55-5 HCAPLUS

CN D-Alaninamide, N-[[[(2S)-tetrahydro-2-furanyl]carbonyl]glycyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-3-(3-pyridinyl)-D-alanyl-N-methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-methylethyl)-L-lysyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 11 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:674362 HCAPLUS

DOCUMENT NUMBER: 142:86565

TITLE: Studies on experimental iodine allergy: 3. Low molecular weight elicitoric antigens of iodine allergy

AUTHOR(S): Sugihara, Yoshiki; Shionoya, Hiroshi; Okano, Kazuo; Sagami, Fumio; Mikami, Takashi; Katayama, Kouichi
CORPORATE SOURCE: Department of Drug Safety Research, Eisai Tsukuba Research Laboratories, Eisai Co., Ltd., Tsukubashi, Ibaraki, 300-2635, Japan

SOURCE: Journal of Toxicological Sciences (2004), 29(2), 147-154

CODEN: JTSCDR; ISSN: 0388-1350

PUBLISHER: Japanese Society of Toxicology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We hypothesize that iodine allergy is an immune response to iodinated self proteins produced in vivo from various iodine-containing chems. Since an antigenic determinant of exptl. iodine allergy is diiodotyrosine (DIT), we designed low mol. weight DIT derivs. having provocative antigenicity without sensitizing immunogenicity. Tetraiododityrosine and hexaiodotrityrosine provoked dose-dependent skin reactions in guinea pigs previously immunized with iodine. No guinea pigs immunized with hexaiodotrityrosine showed anaphylactic reaction by i.v. challenge with hexaiodotrityrosine and none of their antisera showed pos. passive cutaneous anaphylaxis (PCA) reaction in guinea pigs, indicating the non-immunogenic nature of the compound Erythrosine, one of the color additives having a structure common with DIT, was assessed for its immunol. property. ELISA inhibition studies on erythrosine revealed that the inhibitory activity of erythrosine was stronger than that of DIT. Furthermore, erythrosine provoked a PCA reaction in animals sensitized with anti-iodine antisera. In conclusion, hexaiodotrityrosine is thought to be useful for skin testing of iodine allergy without any fear of sensitization to the allergen. Erythrosine was shown to provoke an exptl. iodine allergy and, also, the relationships

09890219

between the new concept of iodine allergy and features of clin. findings of adverse effects by iodocontrast media are discussed.

IT 174608-41-4


RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); BIOL (Biological study)

(low mol. weight DIT derivative hexaiodotriptyrosine is antigenic but not immunogenic in provoking PCA in guinea pig iodine allergy model and erythrosine is immunogenic in provoking exptl. iodine allergy)

RN 174608-41-4 HCAPLUS

CN L-Tyrosine, 3,5-diiodo-L-tyrosyl-3,5-diiodo-L-tyrosyl-3,5-diiodo- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 12 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:466148 HCAPLUS

DOCUMENT NUMBER: 141:174467

TITLE: Total synthesis of eurypamides, marine cyclic-isodityrosines from the Palauan sponge Microciona eurypa

AUTHOR(S): Ito, Miyuki; Yamanaka, Maki; Kutsumura, Noriki; Nishiyama, Shigeru

CORPORATE SOURCE: Department of Chemistry, Faculty of Science and Technology, Keio University, Kohoku-ku, Yokohama, 223-8522, Japan

SOURCE: Tetrahedron (2004), 60(26), 5623-5634

CODEN: TETRAB; ISSN: 0040-4020

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 141:174467

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Total synthesis of eurypamides A, B, and D has been successfully

Updated Search

accomplished. The $\text{Ti}(\text{NO}_3)_3$ (TTN) oxidation of the halogenated bisphenols, e. g. I (Boc = tert-butoxycarbonyl), effected regio-controlled cyclization to provide the corresponding diaryl ethers, e. g. II. This investigation revealed a structural revision of euryamide A as to possess (2''S,3''R,4''S)-configuration, along with the spectral data of pure euryamides A and D, which were previously characterized in a mixture

IT 620960-60-3P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(total synthesis of euryamides, isolated from *Microciconia eurypha*, via $\text{Ti}(\text{NO}_3)_3$ oxidation and regioselective cyclization of halogenated bisphenols intermediates)

RN 620960-60-3 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-3,5-diiodo-L-tyrosyl-L-threonyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 13 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:353144 HCAPLUS

DOCUMENT NUMBER: 140:368700

TITLE: Methods using exemestane, alone or with other therapeutic agents, for treating estrogen-dependent disorders

INVENTOR(S): Wajszczuk, Charles Paul; Gans, Hendrik J. Dekoning; Di Salle, Enrico; Piscitelli, Gabriella; Massimini, Giorgio; Purandare, Dinesh

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 21 pp., Cont.-in-part of WO 2002 72,106.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004082557	A1	20040429	US 2003-611653	20030702
WO 2002072106	A2	20020919	WO 2002-EP638	20020118
WO 2002072106	A3	20031030		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,

09890219

LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
UA, UG, US, UZ, VN, YU, ZA, ZM, ZW
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2001-770911 B2 20010126
WO 2002-EP638 A2 20020118
US 2002-393320P P 20020702

AB The invention discloses a method of preventing and/or treating
estrogen-dependent disorders selected from endometriosis, uterine
fibroids, dysfunctional uterine bleeding, endometrial hyperplasia,
polycystic ovarian disease, fibrocystic breast disease and fibrocystic
mastopathy, which comprises administering to a female mammal in need of
such treatment an effective amount of aromatase inactivator exemestane,
alone or in combination with addnl. therapeutic agents. The invention
also discloses a method for treating infertility in a female mammal in
need of the infertility treatment, comprising administering an effective
amount of exemestane to the mammal.


IT 428438-55-5, A 84861
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL
(Biological study); USES (Uses)
(exemestane, alone or with other therapeutic agents, for treating
estrogen-dependent disorders)

RN 428438-55-5 HCAPLUS


CN D-Alaninamide, N-[[[(2S)-tetrahydro-2-furanyl]carbonyl]glycyl-3-(2-
naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-3-(3-pyridinyl)-D-alanyl-N-
methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-
methylethyl)-L-lysyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



Updated Search



L11 ANSWER 14 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:209417 HCAPLUS

DOCUMENT NUMBER: 141:157439

TITLE: TMC-95A analogues with endocyclic biphenyl ether group as proteasome inhibitors

AUTHOR(S): Kaiser, Markus; Milbradt, Alexander G.; Siciliano, Carlo; Assfalg-Machleidt, Irmgard; Machleidt, Werner; Groll, Michael; Renner, Christian; Moroder, Luis

CORPORATE SOURCE: Max-Planck-Institut fuer Biochemie, AG Bioorganische Chemie, Martinsried, D-82152, Germany

SOURCE: Chemistry & Biodiversity (2004), 1(1), 161-173

CODEN: CBHIAM; ISSN: 1612-1872

PUBLISHER: Verlag Helvetica Chimica Acta AG

DOCUMENT TYPE: Journal

LANGUAGE: English

AB TMC-95A, a cyclic tripeptide metabolite of *Apiospora montagnei*, is a potent competitive inhibitor of proteasome. Based on the X-ray structure of its complex with yeast proteasome, the synthetically challenging structure of this natural product was simplified in a first generation of analogs by replacing the highly oxidized side-chain biaryl system with a phenyl-oxindole group. In the present study, the TMC-95 biaryl group was substituted with a biphenyl ether with retainment of significant proteasome inhibition. Because of the facile synthetic access of tripeptides containing in i, i+2 positions residues of the isodityrosine type, this new generation of TMC-95 analogs may represent promising lead structures for further optimization of affinity and selectivity of proteasome inhibitors.

IT 728007-88-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


(preparation of TMC-95A analogs with endocyclic biphenyl ether group as proteasome inhibitors)

RN 728007-88-3 HCAPLUS

CN L-Tyrosinamide, 3-fluoro-4-nitro-N-[(phenylmethoxy)carbonyl]-L-phenylalanyl-L-asparaginyl-N-propyl-O-[tris(1-methylethyl)silyl]- (9CI)
(CA INDEX NAME)

09890219

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 15 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:989207 HCAPLUS

DOCUMENT NUMBER: 140:287691

TITLE: Development and characterization of potent and specific peptide inhibitors of p60c-src protein tyrosine kinase using pseudosubstrate-based inhibitor design approach

AUTHOR(S): Kamath, J. R.; Liu, R.; Enstrom, A. M.; Lou, Q.; Lam, K. S.

CORPORATE SOURCE: Division of Hematology and Oncology, Department of Internal Medicine, UC Davis Cancer Center, University of California Davis, Sacramento, CA, 95817, USA

SOURCE: Journal of Peptide Research (2003), 62(6), 260-268
CODEN: JPERFA; ISSN: 1397-002X

PUBLISHER: Blackwell Publishing Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 140:287691

AB The cytoplasmic protein p60c-src, an ubiquitous non-receptor protein tyrosine kinase (PTK) is a potential anticancer target as it is over-expressed and/or constitutively active in several cancer types. In addition, the phenotype of c-src knock-out mice is consistent with osteopetrosis, which suggests that inhibitors against this enzyme may also be therapeutic for osteoporosis. Using a known peptide substrate for c-src, MIYKYYF, as a template, we have developed a series of pseudosubstrate-based peptide inhibitors. Structure-activity relationship studies have been performed on one of these inhibitors, CIYKYYF. In a kinase assay using YIYGSFK as the substrate, CIYKYY has been demonstrated to inhibit p60c-src, with an IC₅₀ of 0.6 μ M. Further truncation has led to the determination that even the smaller peptide, CIYK, is a moderately potent inhibitor with IC₅₀ of 15 μ M. Some improvement in inhibitory potency (IC₅₀ = 11.8 μ M) has been observed with the replacement of Tyr3 in CIYK with β -phenylalanine (β -Phe). The tetrapeptide Cl(β -Phe)K will be used as a lead compound for future development of

Updated Search

peptidomimetics and small mol. inhibitors that have the capacity to
penetrate the plasma membrane of intact cells.


IT 676144-47-1P
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation);
BIOL (Biological study); PREP (Preparation)
(development and characterization of peptide inhibitors of p60c-src
protein tyrosine kinase)

RN 676144-47-1 HCAPLUS
CN L-Phenylalaninamide, L-cysteinyl-L-isoleucyl-3,5-diiodo-L-tyrosyl-L-lysyl-
L-tyrosyl-L-tyrosyl- (9CI) (CA INDEX NAME)

[illegible]

L11 ANSWER 16 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2003:769309 HCAPLUS
DOCUMENT NUMBER: 139:365204
TITLE: Synthesis and structural revision of eurypamides
isolated from the Palauan sponge Microciona eurypa
AUTHOR(S): Ito, Miyuki; Yamanaka, Maki; Kutsumura, Noriki;
Nishiyama, Shigeru
CORPORATE SOURCE: Faculty of Science and Technology, Department of
Chemistry, Keio University, Kohoku-ku, Yokohama,
223-8522, Japan
SOURCE: Tetrahedron Letters (2003), 44(43), 7949-7952
CODEN: TELEAY; ISSN: 0040-4039
PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 139:365204
GI

Updated Search




AB Eurypamides A and B were successfully synthesized by employing $\text{Tl}(\text{NO}_3)_3$ (TTN) oxidation of the corresponding halogenated phenols I [$\text{R}_1 = \text{CH}[(\text{R})-\text{Me}]\text{OH}$, $\text{CH}[(\text{S})-\text{OTBS}]\text{CH}[(\text{R})-\text{OTBS}]\text{CH}_2\text{N}_3$, $\text{CH}[(\text{R})-\text{OTBS}]\text{CH}[(\text{S})-\text{OTBS}]\text{CH}_2\text{N}_3$; $\text{TBS} = \text{SiMe}_2\text{t-Bu}$; $\text{Boc} = \text{CO}_2\text{t-Bu}$]. This investigation revealed that the dihydroxyarginine residue of eurypamide A should be revised to possess (2S,3R,4S)-configuration. In addition, the synthesis of eurypamide B provided a pure sample, which was previously characterized in a mixture

IT 620960-60-3P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (synthesis via $\text{Tl}(\text{NO}_3)_3$ oxidation and cyclization of phenols intermediates of eurypamides A and B isolated from *Microciona eurypa*)

RN 620960-60-3 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-3,5-diiodo-L-tyrosyl-L-threonyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 17 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:533623 HCAPLUS

DOCUMENT NUMBER: 139:277151

TITLE: Diastereoselective synthesis of 1-benzyltetrahydroisoquinoline derivatives from amino acids by 1,4 chirality transfer. Part 2


AUTHOR(S): Zawadzka, Anna; Leniewski, Andrzej; Maurin, Jan K.; Wojtasiewicz, Krystyna; Siwicka, Aleksandra; Blachut, Dariusz; Czarnocki, Zbigniew

CORPORATE SOURCE: Faculty of Chemistry, Warsaw University, Warsaw, 02-093, Pol.


SOURCE: European Journal of Organic Chemistry (2003), (13), 2443-2453

PUBLISHER:
DOCUMENT TYPE:
LANGUAGE:
OTHER SOURCE(S):
GI

CODEN: EJOCFK; ISSN: 1434-193X
Wiley-VCH Verlag GmbH & Co. KGaA
Journal
English
CASREACT 139:277151



I



II

AB L-Amino acids (L-Ala, L-Phe, L-Val, L-Pro) were used as a source of chirality in the diastereoselective synthesis of tetrahydroisoquinoline derivs. The key step was the Pictet-Spengler condensation of ketoamides I [R1 = H or Me; R2 is an amino acid side chain; or R1R2 = (CH2)3; R3 = H, Cl], which proceeded under very mild conditions. L-Ala, L-Phe and L-Val gave rise the R-configuration at the newly formed stereogenic center. Surprisingly, L-Pro gave the opposite result. The stereochem. of II [R1, R2 = Me, R4 = Cl; R1R2 = (CH2)3; R3 = H] were established on the basis of X-ray crystallog. data.

IT 603121-75-1P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(diastereoselective synthesis of benzylisoquinoline pyrazinedione derivs. from amino acids via Pictet-Spengler condensation)

RN 603121-75-1 HCAPLUS

CN Benzenepropanamide, 3-chloro-N-[(1S)-2-[[2-(3,4-dimethoxyphenyl)ethyl]amino]-1-methyl-2-oxoethyl]-N-methyl- α -oxo-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT:

38

THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 18 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

Updated Search

09890219

ACCESSION NUMBER: 2003:343939 HCAPLUS
DOCUMENT NUMBER: 139:286537
TITLE: Established theory of radiation-induced decay is not
generalizable to Bolton-Hunter labeled peptides
AUTHOR(S): Doran, Amanda C.; Wan, Yieh-Ping; Kopin, Alan S.;
Beinborn, Martin
CORPORATE SOURCE: Molecular Cardiology Research Institute, Molecular
Pharmacology Research Center, Tufts-New England
Medical Center, Boston, MA, 02111, USA
SOURCE: Biochemical Pharmacology (2003), 65(9), 1515-1520
CODEN: BCPA6; ISSN: 0006-2952
PUBLISHER: Elsevier Science Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English



AB Peptide hormones radiolabeled with ^{125}I are widely used for the pharmacol. characterization of cognate receptors. As a prerequisite for calculating ligand affinities from competition binding assays, and for estimating receptor densities from such studies, it is necessary to know the concentration of bioactive radioligand that is used in resp. expts. It has been demonstrated previously that radioiodinated peptides undergo decay catastrophe, i.e., disintegration of the radioactive label leads to the concomitant destruction of the carrier peptide. Decay catastrophe does not apply to two peptide hormones that are iodinated by Bolton-Hunter conjugation: cholecystokinin octapeptide and glucagon-like peptide 2. The function of aged samples of these radioligands at corresponding recombinantly expressed receptors was assessed by measuring ligand-induced inositol phosphate production or generation of cAMP, resp. Both of the tested compds., although predicted by decay catastrophe to contain little or subthreshold remaining bioactivity, stimulated an unexpectedly high level of receptor-mediated second messenger signaling. Quant. comparison of observed functions with those of corresponding unlabeled peptides suggested that the bioactivity of each radioligand had been largely conserved despite the radioactive decay of the iodine label. Consistent with an apparent absence of decay catastrophe, the authors noted that the specific radioactivity, when determined immediately following peptide iodination, was close to the theor. maximum but exponentially decreased over time. These findings raise the possibility that attachment of a Bolton-Hunter conjugate may shield labeled peptides from radiation-induced damage, a scenario that should be considered when performing radioligand binding expts.

IT 79672-09-6
RL: ANT (Analyte); BSU (Biological study, unclassified); CPS (Chemical process); PEP (Physical, engineering or chemical process); ANST (Analytical study); BIOL (Biological study); PROC (Process)
(established theory of radiation-induced decay is not generalizable to Bolton-Hunter labeled peptides in relation to second messenger signaling in COS-7 cells)

RN 79672-09-6 HCAPLUS

CN Cholecystokinin-8 (swine), N-[3-[4-hydroxy-3-(iodo- ^{125}I)phenyl]-1-oxopropyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 19 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2003:133081 HCAPLUS
DOCUMENT NUMBER: 138:193268

Updated Search

09890219

TITLE: Polypeptide conjugates with extended circulating half-lives
 INVENTOR(S): West, Theodore R.; McMurry, Thomas J.; Dumas, Stephane; Kolodziej, Andrew.
 PATENT ASSIGNEE(S): Epix Medical, Inc., USA
 SOURCE: PCT Int. Appl., 62 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003013573	A1	20030220	WO 2002-US25323	20020809
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2002324655	A1	20030224	AU 2002-324655	20020809
EP 1423136	A1	20040602	EP 2002-759312	20020809
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK			
JP 2004537580	T	20041216	JP 2003-518579	20020809
US 2004254119	A1	20041216	US 2004-487025	20040715
US 7186797	B2	20070306		
JP 2006063071	A	20060309	JP 2005-218098	20050727
PRIORITY APPLN. INFO.:			US 2001-311557P	P 20010810
			JP 2003-518579	A3 20020809
			WO 2002-US25323	W 20020809

OTHER SOURCE(S): MARPAT 138:193268

AB The present invention relates to compds. and methods for synthesizing compds. wherein the compds. exhibit extended circulating half-life in the blood. The increase in circulating half-life is achieved by conjugating polypeptides to binding groups that exhibit high affinity for human serum albumin. A conjugate of recombinant human insulin with 5-oxy-pentanoic acid-phosphono-(R)-2-oxymethyl-N-(S)-4-isobutyl- α -methylphenylacetyl-N'-(S)-N-(3-methoxy-2,4,6-triiodobenzamide)aspartate ethylenediamine diamide was prepared, and its pharmacokinetics was examined in rabbit.


IT 497937-19-6D, derivs., conjugates with polypeptides
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (preparation of polypeptide conjugates having higher affinity for human serum albumin)

RN 497937-19-6 HCAPLUS

CN Benzenepropanamide, N,N'-[(1R)-1-[(phosphonooxy)methyl]-1,2-ethanediyl]bis[2,4,6-triiodo-3-methoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 20 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:716096 HCAPLUS

DOCUMENT NUMBER: 137:226651

TITLE: Combined method for treating hormone-dependent disorders with aromatase inactivator exemestane and other therapeutic agents

INVENTOR(S): Di Salle, Enrico; Piscitelli, Gabriella; Massimini, Giorgio; Purandare, Dinesh; Dekoning, Gans Hendrik

PATENT ASSIGNEE(S): Pharmacia Italia S.p.A., Italy; Pharmacia & Upjohn Company

SOURCE: PCT Int. Appl., 49 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002072106	A2	20020919	WO 2002-EP638	20020118
WO 2002072106	A3	20031030		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2434611	A1	20020919	CA 2002-2434611	20020118
AU 2002257573	A1	20020924	AU 2002-257573	20020118
EP 1377298	A2	20040107	EP 2002-727314	20020118
EP 1377298	B1	20060830		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2004519490	T	20040702	JP 2002-571065	20020118
AT 337787	T	20060915	AT 2002-727314	20020118
US 2004082557	A1	20040429	US 2003-611653	20030702
PRIORITY APPLN. INFO.:				
			US 2001-770911	A 20010126
			WO 2002-EP638	W 20020118

AB A method of preventing and treating estrogen dependent disorders selected from endometriosis, uterine fibroids, dysfunctional uterine bleeding, endometrial hyperplasia, polycystic ovarian disease, fibrocystic breast disease and fibrocystic mastopathy, is disclosed which is comprised of administering to a mammalian patient in need of such treatment an effective amount of aromatase inactivator exemestane, alone or in combination with addnl. therapeutic agents.

IT 428438-55-5, A 84861

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(combined method for treating hormone-dependent disorders with aromatase inactivator exemestane and other therapeutic agents)


RN 428438-55-5 HCAPLUS

CN D-Alaninamide, N-[[[(2S)-tetrahydro-2-furanyl]carbonyl]glycyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-3-(3-pyridinyl)-D-alanyl-N-methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-methylethyl)-L-lysyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A






PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002070511	A1	20020912	WO 2002-US6479	20020302
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2437594	A1	20020912	CA 2002-2437594	20020302
AU 2002254095	A1	20020919	AU 2002-254095	20020302
EP 1363898	A1	20031126	EP 2002-723310	20020302
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
HU 200401544	A2	20041228	HU 2004-1544	20020302
JP 2005511475	T	20050428	JP 2002-569831	20020302
US 2003092732	A1	20030515	US 2002-90582	20020304
US 6979691	B2	20051227		

09890219

US 2003096827	A1	20030522	US 2002-90288	20020304
US 6713487	B2	20040330		
US 2004229882	A1	20041118	US 2003-696761	20031029
US 7067525	B2	20060627		
US 2006025403	A1	20060202	US 2005-199464	20050808
PRIORITY APPLN. INFO.:			US 2001-273206P	P 20010302
			US 2001-273291P	P 20010302
			WO 2002-US6479	W 20020302
			US 2002-90288	A3 20020304
			US 2002-90582	A3 20020304

OTHER SOURCE(S): MARPAT 137:232913
GI



AB Compds. W-(CR6R7)yCH(G)(CR4R5)xCO-X(R1)CHR2(CHR3)r(CH2)sCO-E [X = N or CH; R1, R3 = H or alkyl; R2 = H, aryl, cycloalkyl, heteroaryl, heterocyclyl, (un)substituted alkyl or alkenyl; R1 together with R2 or R3 or R2 together with R3 form mono- or bicyclic aryl, cycloalkyl, heteroaryl, or heterocyclyl; E = (un)substituted pyrrolidino, piperidino, hexahydro-1-azepinyl, 1-piperazinyl, cyclopentyl, cyclohexyl, cycloheptyl, amino, (cyclo)alkylamino; R4-R6 = H, (un)substituted alkyl, amino, alkylamino, hydroxy, alkoxy, aryl, cycloalkyl, heteroaryl, or heterocyclyl; or CR4R5 or C6R7 is a spirocycloalkyl ring; r, s = 0 or 1; x = 0-4; y = 0-2; G = alkenyl, arylalkenyl, hydroxy, heteroaryl, cyano, functionalized alkyl or alkenyl, etc.; W = amino, alkylamino, hydroxy, alkoxy, carbamoyl, amidino, cycloalkyl, heteroaryl, heterocyclyl, etc.] were prepared as modulators of melanocortin receptors, particularly MC-1R and MC-4R. Thus, peptide I was prepared by a solution-phase peptide coupling/deprotection scheme.

IT 457902-69-1P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(preparation of peptides for pharmaceutical use as modulators of melanocortin receptors)

RN 457902-69-1 HCAPLUS

CN 4-Piperidinepropanamide, α -[[[3-(2-chlorophenyl)-1-oxopropyl]amino]-N-[(1R)-1-[(4-methoxyphenyl)methyl]-2-oxo-2-[4-(1-oxobutyl)-4-phenyl-1-piperidinyl]ethyl]-, (α S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 22 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:692478 HCAPLUS

DOCUMENT NUMBER: 138:304504

TITLE: Development and characterization of potent peptide inhibitors of p60c-src PTK using pseudosubstrate-based inhibitor design approach

AUTHOR(S): Kamath, Jayesh R.; Liu, Ruiwu; Enstrom, Amanda M.; Liu, Gang; Lou, Qiang; Lam, Kit S.

CORPORATE SOURCE: Division of Hematology and Oncology, Department of Internal Medicine, UC Davis Cancer Center, University of California Davis, Sacramento, CA, 95817, USA

SOURCE: Peptides: The Wave of the Future, Proceedings of the Second International and the Seventeenth American Peptide Symposium, San Diego, CA, United States, June 9-14, 2001 (2001), 551-552. Editor(s): Lebl, Michal; Houghten, Richard A. American Peptide Society: San Diego, Calif.

CODEN: 69DBAL; ISBN: 0-9715560-0-8

DOCUMENT TYPE: Conference

LANGUAGE: English

AB A symposium report. Using MIYKYF as template, potent pseudosubstrate-based inhibitors of p60c-src protein tyrosine kinase were synthesized. SAR (structure-activity relationship) studies of the parent peptide MIYKYF identified IYKYF as an inhibitor with similar potency as its parent peptide. Inhibition evaluations using dithiolthreitol showed a ten-fold reduction in inhibitory potency of CIYKYF, suggesting that disulfide bond formation between the Cys1 of the peptide CIYKYF and a cysteine residue at the enzyme active site could account for the enhanced potency of CIYKYF. At the N-terminus, the sulfhydryl group, the free N α -group and the L-enantiomer of Cys1 are crucial for the improved inhibitory potency of the peptide CIYKYF. The replacement of L-isoleucine2 of CIYKYF with unnatural amino acid D-propargylglycine resulted in the development of the most potent inhibitor of this series. No improvement in inhibitory potency was noted with the Tyr3 mimetic analogs of CIYKYF or CIYK, indicating the possibility that the hydroxyl group of Tyr3 could be essential for certain critical interactions at the enzyme active site.

IT 509149-13-7P

RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)


09890219.

```
(preparation of pseudosubstrate-based potent peptide inhibitors of protein
tyrosine kinase and enzyme-inhibiting structure-activity relationship)
```

RN 509149-13-7 HCAPLUS

CN L-Phenylalanine, L-cysteinyl-L-isoleucyl-3,5-diiodo-L-phenylalanyl-L-lysyl-L-tyrosyl-L-tyrosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 23 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:637480 HCAPLUS

DOCUMENT NUMBER: 137:190724

TITLE: Melanocortin metallopeptides for treatment of sexual dysfunction

INVENTOR(S): Sharma, Shubh D.; Shi, Yi-qun; Yang, Wei; Cai, Hui-zhi; Shadiack, Annette

PATENT ASSIGNEE(S): Palatin Technologies, Inc., USA

SOURCE: PCT Int. Appl., 58 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 6

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002064091	A2	20020822	WO 2002-US4431	20020213
WO 2002064091	A3	20030313		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2002238106	A1	20020828	AU 2002-238106	20020213
US 2004038897	A1	20040226	US 2003-640755	20030813
US 2005164914	A1	20050728	US 2005-36273	20050114
PRIORITY APPLN. INFO.:			US 2001-268591P	P 20010213

Updated Search

09890219

US 1995-476652	A2	19950607
US 1996-660697	A3	19960605
US 2000-483837	A2	20000117
WO 2002-US4431	W	20020213
US 2003-640755	A2	20030813
US 2004-536691P	P	20040114

OTHER SOURCE(S) : MARPAT 137:190724

AB Metallopeptides are provided for use in treatment of sexual dysfunction in mammals. The metallopeptides are agonists for at least one of melanocortin-3 or melanocortin-4 receptors. The metallopeptides are conformationally fixed on complexation of a metal ion-binding portion thereof with a metal ion. Also provided are metallopeptides that are antagonists for at least one of melanocortin-3 or melanocortin-4 receptors.


IT 448902-31-6

RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(melanocortin metallopeptides for treatment of sexual dysfunction)

RN 448902-31-6 HCAPLUS

CN L-Cysteinamide, N-(1-oxoheptyl)-O-(phenylmethyl)-L-seryl-4-chloro-D-phenylalanyl-L-arginyl-L-tyrosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 24 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:575744 HCAPLUS

DOCUMENT NUMBER: 137:135069

TITLE: Method for reducing or preventing the establishment, growth or metastasis of cancer by administering indole peptidomimetics PAR-1 antagonist and optionally PAR-2 antagonists

INVENTOR(S): D'Andrea, Michael; Derian, Claudia; Woodrow, Hal Brent
PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 44 pp., Cont.-in-part of U.S. Ser. No. 603,231.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

Updated Search

09890219

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002103138	A1	20020801	US 2001-865824	20010525
US 6858577	B1	20050222	US 2000-603231	20000626
US 2003224999	A1	20031204	US 2003-403542	20030331
US 7183252	B2	20070227		

PRIORITY APPLN. INFO.: US 1999-141550P P 19990629
US 2000-603231 A2 20000626

OTHER SOURCE(S): MARPAT 137:135069

AB The authors have discovered a method of modifying the tumor cell microenvironment to reduce or prevent the establishment, growth or metastasis of malignant cells comprising administering to a patient having malignant cells a pharmaceutically effective amount of a PAR-1 (proteinase-activated receptor 1) inhibitor and optionally a PAR-2 (proteinase-activated receptor 2) inhibitor to prevent or reduce activation of normal cells within the tumor microenvironment. This method also has the effect in some patients of modulating the immune system to facilitate a more efficient immune response to malignant cells and maybe coupled with cytokine therapy and T-cell therapy to enhance the patient's immune response to the malignant cells.


IT 316150-72-8P, L-Argininamide, 3,4-difluoro-N-[[[1-[(2-methylphenyl)methyl]-3-(1-pyrrolidinylmethyl)-1H-indol-6-yl]amino]carbonyl]-L-phenylalanyl-N-[2-(2-methoxyphenyl)ethyl]-
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(inhibition of growth or metastasis of cancer by administering indole peptidomimetics PAR-1 antagonists and combined with PAR-2 antagonists and other agents in relation to immunostimulant activity)

RN 316150-72-8 HCAPLUS

CN L-Argininamide, 3,4-difluoro-N-[[[1-[(2-methylphenyl)methyl]-3-(1-pyrrolidinylmethyl)-1H-indol-6-yl]amino]carbonyl]-L-phenylalanyl-N-[2-(2-methoxyphenyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 25 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:487180 HCAPLUS

DOCUMENT NUMBER: 137:228443

TITLE: A CLN2-related and thermostable serine-carboxyl

Updated Search

proteinase, kumamolysin: cloning, expression, and identification of catalytic serine residue

AUTHOR(S): Oyama, Hiroshi; Hamada, Takatoshi; Ogasawara, Shin; Uchida, Kenichi; Murao, Sawao; Beyer, Bret B.; Dunn, Ben M.; Oda, Kohei

CORPORATE SOURCE: Department of Applied Biology, Faculty of Textile Science, Kyoto Institute of Technology, Kyoto, 606-8585, Japan

SOURCE: Journal of Biochemistry (Tokyo, Japan) (2002), 131(5), 757-765
CODEN: JOBIAO; ISSN: 0021-924X

PUBLISHER: Japanese Biochemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The gene encoding kumamolysin, a thermostable pepstatin-insensitive carboxyl proteinase, was cloned and expressed. (i) Kumamolysin was synthesized as a large precursor consisting of two regions: amino-terminal prepro (188 amino acids) and mature proteins (384 amino acids). (ii) The deduced amino acid sequence of the mature region exhibited high similarity to those of such bacterial pepstatin-insensitive enzymes as *Pseudomonas* carboxyl proteinase (PSCP; EC 3.4.23.37, identity = 37%), *Xanthomonas* carboxyl proteinase (XCP; EC 3.4.23.33, identity = 36%), and human CLN2 gene product (identity = 36%), which is related to a fatal neurodegenerative disease. (iii) The presumed catalytic triad, Glu78, Asp82, Ser278, was found to be conserved in the amino acid sequence of kumamolysin. (iv) Kumamolysin was inactivated by such aldehyde-type inhibitors as Ac-Ile-Pro-Phe-CHO ($K_i = 0.7 \pm 0.14 \mu\text{M}$): In PSCP, it has been clarified that these inhibitors form a hemiacetal linkage with the catalytic serine residue and inactivate the enzyme. (v) Mutational anal. of the Ser278 residue revealed that the mutant lost both auto-processing activity and proteolytic activity. These results strongly suggest that kumamolysin has a unique catalytic triad consisting of Glu78, Asp82, and Ser278 residues, as previously observed for PSCP.


IT 392232-98-3
RL: ARU (Analytical role, unclassified); BSU (Biological study, unclassified); PRP (Properties); ANST (Analytical study); BIOL (Biological study)

(kumamolysin has a unique catalytic triad consisting of Glu78, Asp82, and Ser278 residues, as previously observed for PSCP)

RN 392232-98-3 HCAPLUS

CN L-Leucinamide, 4-iodo-N-(3-methyl-1-oxobutyl)-L-phenylalanyl-N-[(1S)-1-formyl-2-(4-hydroxyphenyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

09890219

L11 ANSWER 26 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:391511 HCAPLUS

DOCUMENT NUMBER: 136:406856

TITLE: Combined therapy against tumors comprising estramustine phosphate and LHRH agonists or antagonists

INVENTOR(S): Buchalter, Jeffrey H.; Horak, Ivan D.

PATENT ASSIGNEE(S): Pharmacia & Upjohn Company, USA

SOURCE: PCT Int. Appl., 11 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent



LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2002039996	A2	20020523	WO 2001-US44161	20011106
WO 2002039996	A3	20030320		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2002028648	A5	20020527	AU 2002-28648	20011106
PRIORITY APPLN. INFO.:			US 2000-714606	A1 20001116
			WO 2001-US44161	W 20011106
AB	A method for treating tumors in a mammal, including humans, in need of such a treatment including administering simultaneously, sep. or sequentially to said mammal estramustine phosphate and a LHRH agonist or antagonist, in amts. sufficient to achieve a therapeutically useful effect. Estramustine phosphate arginine salt formulation for injection was prepared			
IT	428438-55-5, A 84861			
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (combined therapy against tumors comprising estramustine phosphate and LHRH agonists or antagonists)			
RN	428438-55-5 HCAPLUS			
CN	D-Alaninamide, N-[[[(2S)-tetrahydro-2-furanyl]carbonyl]glycyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-3-(3-pyridinyl)-D-alanyl-N-methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-methylethyl)-L-lysyl-L-prolyl- (9CI) (CA INDEX NAME)			

Absolute stereochemistry.



L11 ANSWER 27 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:391510 HCAPLUS

DOCUMENT NUMBER: 136:380114

TITLE: Aromatase inhibitor combination with inhibition of testicular and ovarian hormone output for treatment of estrogen-dependent cancers

INVENTOR(S): Purandare, Dinesh
 PATENT ASSIGNEE(S): Pharmacia & Upjohn Company, USA
 SOURCE: PCT Int. Appl., 13 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1



PATENT INFORMATION:

09890219

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002039995	A2	20020523	WO 2001-US43847	20011106
WO 2002039995	A9	20030206		
WO 2002039995	A3	20030501		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2428249	A1	20020523	CA 2001-2428249	20011106
AU 200230464	A	20020527	AU 2002-30464	20011106
EP 1341549	A2	20030910	EP 2001-990699	20011106
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
CN 1498112	A	20040519	CN 2001-818938	20011106
JP 2004536022	T	20041202	JP 2002-542370	20011106
BR 2001015423	A	20051213	BR 2001-15423	20011106
NZ 525720	A	20061222	NZ 2001-525720	20011106
ZA 2003003669	A	20040513	ZA 2003-3669	20030513
NO 2003002206	A	20030715	NO 2003-2206	20030515
US 2004043938	A1	20040304	US 2003-416844	20030912
PRIORITY APPLN. INFO.:			US 2000-714605	A1 20001116
			WO 2001-US43847	W 20011106
AB	The invention provides a combination therapy for treating estrogen-dependent cancers in susceptible mammals, including humans, comprising inhibiting testicular or ovarian hormone output and administering at least one aromatase inhibitor.			
IT	428438-55-5, A 84861 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (aromatase inhibitor combination with inhibition of testicular and ovarian hormone output for treatment of estrogen-dependent cancers)			
RN	428438-55-5 HCAPLUS			
CN	D-Alaninamide, N-[[[(2S)-tetrahydro-2-furanyl]carbonyl]glycyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-3-(3-pyridinyl)-D-alanyl-N-methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-methylethyl)-L-lysyl-L-prolyl- (9CI) (CA INDEX NAME)			

Absolute stereochemistry.

Updated Search



L11 ANSWER 28 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001:857934 HCAPLUS
 DOCUMENT NUMBER: 136:147059
 TITLE: Inhibitor Complexes of the Pseudomonas Serine-Carboxyl Proteinase
 AUTHOR(S): Wlodawer, Alexander; Li, Mi; Gustchina, Alla; Dauter, Zbigniew; Uchida, Kenichi; Oyama, Hiroshi; Goldfarb, Nathan E.; Dunn, Ben M.; Oda, Kohei
 CORPORATE SOURCE: Protein Structure Section Macromolecular Crystallography Laboratory and Intramural Research Support Program, SAIC Frederick National Cancer Institute at Frederick, Frederick, MD, 21702, USA
 SOURCE: Biochemistry (2001), 40(51), 15602-15611
 CODEN: BICHAW; ISSN: 0006-2960
 PUBLISHER: American Chemical Society

09890219

DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 136:147059


AB Crystal structures of the serine-carboxyl proteinase from *Pseudomonas* sp. 101 (PSCP), complexed with a number of inhibitors, have been solved and refined at high- to atomic-level resolution. All of these inhibitors (tyrostatin, pseudo-tyrostatin, AcIPF, AcIAF, and chymostatin, as well as previously studied iodotyrostatin and pseudo-iodotyrostatin) make covalent bonds to the active site Ser287 through their aldehyde moieties, while their side chains occupy subsites S1-S4 of the enzyme. The mode of binding of the inhibitors is almost identical for their P1 and P2 side chains, while significant differences are observed for P3 and P4 (if present). Kinetic parameters for the binding of these nanomolar inhibitors to PSCP have been established and correlated with the observed mode of binding. The preferences of this enzyme for a larger side chain in P2 as well as Tyr or Phe in P1 are explained by the size, shape, and characteristics of the S2 and S1 regions of the protein structure, resp. Networks of hydrogen bonds involving glutamic and aspartic acids have been analyzed for the atomic-resolution structure of the native enzyme. PSCP contains a calcium-binding site that consists of Asp328, Asp348, three amide carbonyl groups, and a water mol., in almost perfect octahedral coordination. The presence of Ca²⁺ cation is necessary for the activity of the enzyme.

IT 392232-98-3P
RL: BSU (Biological study, unclassified); SPN (Synthetic preparation);
BIOL (Biological study); PREP (Preparation)
(preparation of peptide inhibitors of the *Pseudomonas* serine-carboxyl proteinase and crystallog. study of the proteinase-inhibitor complexes)

RN 392232-98-3 HCAPLUS

CN L-Leucinamide, 4-iodo-N-(3-methyl-1-oxobutyl)-L-phenylalanyl-N-[(1S)-1-formyl-2-(4-hydroxyphenyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 29 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:615130 HCAPLUS

DOCUMENT NUMBER: 135:358138

TITLE: Synthesis of a cyclic diaryl ether derivative under solid-phase conditions

AUTHOR(S): Nakamura, K.; Nishiya, H.; Nishiyama, S.

CORPORATE SOURCE: National Institute of Advanced Industrial Science and Technology, Tsukuba, 305-8566, Japan

SOURCE: Tetrahedron Letters (2001), 42(36), 6311-6313

CODEN: TELEAY; ISSN: 0040-4039

PUBLISHER: Elsevier Science Ltd.

Updated Search

09890219

DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 135:358138

AB The TTN phenolic oxidation, along with the N-protective group of the corresponding tripeptide derivs., was examined to accomplish construction of a cyclic isodityrosine derivative under solid-phase conditions. The desired cyclization was effected under the TTN (thallium(III) trinitrate)/NMP-MeOH conditions to give the corresponding 17-membered ring lactam 12.

IT 372963-97-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of cyclic isodityrosine derivative from tripeptides by phenolic oxidation using TTN under solid-phase conditions)

RN 372963-97-8 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-3,5-diiodo-L-tyrosyl-L-isoleucyl-3,5-dibromo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 30 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:325496 HCAPLUS

DOCUMENT NUMBER: 136:17330

TITLE: Antitumor and antiangiogenic effects of somatostatin receptor-targeted in situ radiation with ¹¹¹In-DTPA-JIC 2DL

AUTHOR(S): Gulec, Seza A.; Drouant, George J.; Fuselier, Joseph; Anthony, Catherine T.; Heneghan, James; DelCarpio, Joseph B.; Coy, David H.; Murphy, William A.; Woltering, Eugene A.

CORPORATE SOURCE: Department of Surgery, The Louisiana State University Health Sciences Center, New Orleans, LA, USA

SOURCE: Journal of Surgical Research (2001), 97(2), 131-137
CODEN: JSGRA2; ISSN: 0022-4804

PUBLISHER: Academic Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Expression of somatostatin receptor subtype 2 (sst 2) in angiogenic tumor vessels appears to be homogeneous, while tumor cell expression of this receptor is often heterogeneous. We have developed a novel in vitro three-dimensional tumor angiogenesis model to study the antitumor and the antiangiogenic effects of radiolabeled somatostatin analogs. We hypothesized that targeted in situ radiation with an Auger electron-emitting radiolabeled somatostatin analog would produce receptor-specific cytotoxicity in sst 2-expressing cells. IMR-32 human

Updated Search

neuroblastoma (sst 2-pos.) and MDA MB-231 human breast cancer (sst 2-neg.) xenografts were created in nude mice from monolayer cell cultures. Fragments of these tumors were embedded in three-dimensional fibrin gels supplemented with endothelial growth media and incubated for a period of 14 days. Tumor fragments were treated with 50 $\mu\text{Ci/mL}$ of ^{111}In -JIC 2DL, a sst 2-preferring somatostatin analog, or medium on Day 1. Initial angiogenic activity was determined at 48 h and the mean angiogenic score and tumoricidal responses were assessed on Day 14. Results and conclusion. Tumoricidal effects of ^{111}In -JIC 2DL were seen only in sst 2-pos. IMR-32 tumors. However, the angiogenic response was inhibited in both IMR-32 and MDA MB-231 tumors independent of the tumor cells sst 2 status. Somatostatin receptor-mediated in situ radiation therapy has profound cytotoxic effects on angiogenic blood vessels and sst 2-expressing tumor cells. (c) 2001 Academic Press.

IT 271785-12-7D, indium-111 DTPA derivative


RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(antitumor and antiangiogenic effects of somatostatin receptor-targeted in situ radiation with ^{111}In -DTPA-JIC 2DL)

RN 271785-12-7 HCAPLUS

CN L-Threoninamide, D-lysyl-3,5-diiodo-D-tyrosyl-D-lysyl-3,5-diiodo-D-tyrosyl-D-lysyl-L-cysteinyl-L-phenylalanyl-D-tryptophyl-L-lysyl-L-threonyl-L-cysteinyl-, cyclic (6-11)-disulfide (9CI) (CA INDEX NAME)

PAGE 1-A





REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 31 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2001:12482 HCAPLUS

DOCUMENT NUMBER: 134:71906

TITLE: Preparation of novel indole peptidomimetics as thrombin receptor antagonists

INVENTOR(S): Zhang, Han-cheng; Hoekstra, William J.; Maryanoff, Bruce E.; McComsey, David F.

PATENT ASSIGNEE(S): Ortho-Mcneil Pharmaceutical, Inc., USA; Cor Therapeutics, Inc.

SOURCE: PCT Int. Appl., 76 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001000657	A2	20010104	WO 2000-US18018	20000629
WO 2001000657	A3	20010712		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 6858577	B1	20050222	US 2000-603231	20000626
US 2003224999	A1	20031204	US 2003-403542	20030331
US 7183252	B2	20070227		

PRIORITY APPLN. INFO.: US 1999-141550P P 19990629

US 2000-603231 A 20000626

OTHER SOURCE(S): MARPAT 134:71906

09890219

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


AB Indole derivs. I [A1 and A2 are certain D- or L-amino acid residues which may be substituted; R1 = amino, alkylamino, arylamino, heteroalkyl, etc.; R2 = H, halo, alkyl, cycloalkyl, alkenyl, alkynyl, arylalkyl, aryl, heteroaryl; R3, R4 = H, alkyl, cycloalkyl, cycloalkylalkyl, aryl, heteroalkyl, indanyl, etc. or R3R4N = (un)substituted piperidinyl, piperazinyl, morpholino, or pyrrolidinyl; R5 = (un)substituted aryl, arylalkyl, cycloalkyl, heteroaryl; R6 = H, alkyl; X = O, S; m = 0-3; n = 1 or 2; p = 0 or 1] were prepared as thrombin receptor antagonists for the treatment of diseases associated with thrombosis, restenosis, hypertension, heart failure, arrhythmia, inflammation, angina, stroke, atherosclerosis, ischemic conditions, angiogenesis related disorders, cancer, and neurodegenerative disorders. Thus, compound II, prepared by a multistep procedure starting from 6-nitroindole (scheme given), showed IC50 = 0.28 and 0.47 μ M, resp., in the thrombin-induced gel-filtered platelet aggregation and thrombin receptor binding assays.

IT 316150-72-8P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of novel indole peptidomimetics as thrombin receptor antagonists)

RN 316150-72-8 HCAPLUS

CN L-Argininamide, 3,4-difluoro-N-[[[1-[(2-methylphenyl)methyl]-3-(1-pyrrolidinylmethyl)-1H-indol-6-yl]amino]carbonyl]-L-phenylalanyl-N-[2-(2-methoxyphenyl)ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 32 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:910888 HCAPLUS

DOCUMENT NUMBER: 136:290900

TITLE: Substrate specificity and inhibition studies of human serotonin N-acetyltransferase. [Erratum to document

Updated Search

09890219

cited in CA133:39768]

AUTHOR(S): Ferry, Gilles; Loynel, Armelle; Kucharczyk, Nathalie; Bertin, Sophie; Rodriguez, Marianne; Delagrangé, Philippe; Galizzi, Jean-Pierre; Jacoby, Edgar; Volland, Jean-Paul; Lesieur, Daniel; Renard, Pierre; Canet, Emmanuel; Fauchère, Jean-Luc; Boutin, Jean A.

CORPORATE SOURCE: Division de Pharmacologie Moléculaire et Cellulaire, Institut de Recherches Servier, Croissy sur Seine, 78290, Fr.

SOURCE: Journal of Biological Chemistry (2000), 275(50), 39799
CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular Biology

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Throughout the text, all of the Vmax units should read "nmol/min/mg protein" rather than "μmol/min/mg protein". This correction is especially evident on page 8797, right column, next to last sentence in the last paragraph; page 8799, left column, 11th line from the bottom of the page; page 8799, right column, 10th through 14th lines; and in the unit headings of Tables II and III (pages 8799 and 8800, resp.).

IT 274918-26-2
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(substrate specificity and inhibition studies of human serotonin N-acetyltransferase (Erratum))

RN 274918-26-2 HCAPLUS

CN L-Tyrosine, 3-benzo[b]thien-3-yl-L-alanyl-4-fluoro-L-phenylalanyl-3-cyclohexyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 33 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:720707 HCAPLUS

DOCUMENT NUMBER: 134:29690

TITLE: A Novel Synthesis of Biaryl-Containing Macrocycles by a Domino Miyaura Arylboronate Formation: Intramolecular Suzuki Reaction

AUTHOR(S): Carbonnelle, Anny-Claude; Zhu, Jieping


CORPORATE SOURCE: Institut de Chimie des Substances Naturelles, CNRS, Gif-sur-Yvette, 91198, Fr.

SOURCE: Organic Letters (2000), 2(22), 3477-3480

Updated Search

09890219

PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 134:29690
GI



AB A novel macrocyclization procedure is developed on the basis of a domino process. Thus, treatment of linear diiodide I under defined conditions gave a 15-membered m,m-cyclophane via aryl-aryl bond formation. Two distinct cross-coupling manifolds, Miyaura's arylboronic ester synthesis and intramol. Suzuki reaction, proceed in an ordered fashion. Concentration is an important factor for the success of this process.

IT 312493-98-4


RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of biaryl-containing macrocycles by a domino Miyaura arylboronate formation:intramol. Suzuki reaction)

RN 312493-98-4 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-3-iodo-L-tyrosyl-L-leucyl-3-iodo-O-methyl-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 34 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:442999 HCAPLUS

DOCUMENT NUMBER: 133:223020

TITLE: Design and Synthesis of Potent Hexapeptide and Heptapeptide Gonadotropin-Releasing Hormone Antagonists by Truncation of a Decapeptide Analogue Sequence

AUTHOR(S): Yahalom, Dror; Rahimipour, Shai; Koch, Yitzhak; Ben-Aroya, Nurit; Fridkin, Mati

Updated Search

09890219

CORPORATE SOURCE: Departments of Organic Chemistry and Neurobiology,
Weizmann Institute of Science, Rehovot, 76100, Israel
SOURCE: Journal of Medicinal Chemistry (2000), 43(15),
2831-2836
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB A novel strategy for designing reduced-size analogs of the decapeptide gonadotropin-releasing hormone (GnRH) was developed. As opposed to previous attempts to delete residues from either of the peptide's termini, our approach is based upon the known importance of both C- and N-terminals of GnRH analogs for receptor recognition, whereas the central part of the mol. is replaced by a short spacer. The present truncation strategy was successful for generation of reduced-size hexapeptide and heptapeptide antagonists possessing potent antagonistic capacity. The same methodol. was not suitable for the generation of reduced-size agonists, suggesting different conformational characteristics for GnRH agonists and antagonists. A heptapeptide antagonist designed by this method was shown to inhibit serum levels of LH in castrated rats in vivo. Structure-activity studies suggested that the structural preferences for GnRH receptor recognition are similar to those reported for decapeptide antagonists. Our studies resulted in a heptapeptide GnRH antagonist (Ac-D-Nal2-D-Cpa-D-Pal-Gly-Arg-Pro-D-Ala-NH2) with high receptor-binding affinity (IC50 = 7 nM), as compared to that of GnRH itself. The highest affinity of a new hexapeptide antagonist was somewhat lower (IC50 = 45 nM).

IT 292141-37-8P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(design and synthesis of hexa- and heptapeptide gonadotropin-releasing hormone antagonists)


RN 292141-37-8 HCAPLUS

CN D-Alaninamide, N-acetyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-3-(2-pyridinyl)-D-alanyl-D-tyrosyl-L-arginyl-L-prolyl- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L11 ANSWER 35 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:431263 HCAPLUS
 DOCUMENT NUMBER: 133:223013
 TITLE: An efficient total synthesis of K-13, a non-competitive inhibitor of ACE I
 AUTHOR(S): Bigot, Antony; Bois-Choussy, Michele; Zhu, Jieping
 CORPORATE SOURCE: Institut de Chimie des Substances Naturelles, CNRS, Gif-sur-Yvette, 91198, Fr.
 SOURCE: Tetrahedron Letters (2000), 41(23), 4573-4577
 CODEN: TELEAY; ISSN: 0040-4039
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 133:223013
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB An efficient synthesis of K-13 (I), a non-competitive inhibitor of ACE I with an endo biaryl ether bond, is described. The key cycloetherification reaction of linear tripeptide II gave a 17-membered macrocycle in quant. yield.
 IT 291781-71-0
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (total synthesis of the 17-membered cyclopeptide K-13 as a non-competitive inhibitor of ACE I)
 RN 291781-71-0 HCAPLUS
 CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-4-fluoro-3-nitro-L-phenylalanyl-O-methyl-L-tyrosyl-3-hydroxy-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 36 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:368415 HCAPLUS
 DOCUMENT NUMBER: 133:13158
 TITLE: Hydrophilic somatostatin analogs
 INVENTOR(S): Coy, David H.; Murphy, William A.; Woltering, Eugene A.; Fuselier, Joseph A.; Drouant, George
 PATENT ASSIGNEE(S): Tulane University, USA
 SOURCE: PCT Int. Appl., 35 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000031122	A1	20000602	WO 1999-US24532	19991020
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
US 6465613	B1	20021015	US 1998-196259	19981119
CA 2351944	A1	20000602	CA 1999-2351944	19991020
EP 1131343	A1	20010912	EP 1999-955077	19991020
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
JP 2003517999	T	20030603	JP 2000-583949	19991020
PRIORITY APPLN. INFO.:			US 1998-196259	A 19981119
			WO 1999-US24532	W 19991020

OTHER SOURCE(S): MARPAT 133:13158

AB The invention features novel somatostatin analogs that may be readily labeled with toxic or non-toxic detectable labels. These unlabeled and labeled analogs are useful for specifically targeting somatostatin receptor bearing cells, in particular neoplastic cells. Labeled analogs are useful, for example, for tumor localization and detection. Where labeled with a toxic label (e.g., radioactivity), the analogs are useful

09890219

for the targeted delivery of toxicity to somatostatin receptor-bearing cells, in particular neoplastic cells. Also disclosed are methods for treating and detecting neoplasms, and methods for imaging somatostatin receptor-bearing cells.

IT 271785-11-6P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(preparation of labeled or unlabeled hydrophilic somatostatin analogs for targeting somatostatin receptor-bearing cells)

RN 271785-11-6 HCAPLUS

CN L-Threoninamide, D-lysyl-3,5-diiodo-L-tyrosyl-L-lysyl-3,5-diiodo-L-tyrosyl-L-lysyl-L-cysteinyl-L-phenylalanyl-D-tryptophyl-L-lysyl-L-threonyl-L-cysteinyl-, cyclic (6-11)-disulfide (9CI) (CA INDEX NAME)

PAGE 1-A





REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 37 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:283962 HCAPLUS

DOCUMENT NUMBER: 132:304929

TITLE: Method of making mammalian kringle 5 peptide fragments with angiogenesis inhibitory effect by elastase proteolytic cleavage of plasminogen

INVENTOR(S): Davidson, Donald J.

PATENT ASSIGNEE(S): Abbott Laboratories, USA

SOURCE: U.S., 48 pp., Cont.-in-part of U.S. Ser. No. 832,087.

CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6057122	A	20000502	US 1997-851350	19970505
US 5801146	A	19980901	US 1996-643219	19960503
US 5981484	A	19991109	US 1997-832087	19970403
US 6699838	B1	20040302	US 1997-924287	19970905
US 2004138127	A1	20040715	US 2004-753646	20040108
PRIORITY APPLN. INFO.:			US 1996-643219	A2 19960503
			US 1997-832087	A2 19970403
			US 1997-851350	A2 19970505
			US 1997-924287	A1 19970905

AB A method of making mammalian kringle 5 peptide fragments corresponding to the 5th kringle domain of mammalian plasminogen and having angiogenic inhibitory effect is claimed. The method comprises exposing a mammalian plasminogen to elastase at a ratio of about 1:100 to 1:300 (weight/weight) and isolating kringle 5 fragments from the mixture. Kringle 5 peptide fragments were prepared either by porcine elastase proteolytic cleavage of Lys plasminogen or synthesized by standard solid phase Fmoc chemical. The inhibition

09890219

of bovine capillary endothelial cell proliferation and migration by kringle 5 peptide fragments was both potent and specific to the endothelial cells but not normal or tumor cells. Kringle 5 peptide fragments were also produced recombinantly in *Pichia pastoris* and *E. coli*.

IT 199664-87-4P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)


(synthesis of, antiangiogenic kringle 5 peptide; method of making mammalian kringle 5 peptide fragments with angiogenesis inhibitory effect by elastase proteolytic cleavage of plasminogen)

RN 199664-87-4 HCAPLUS


CN L-Tyrosinamide, 1-acetyl-L-prolyl-L-arginyl-L-lysyl-L-leucyl-3-iodo-L-tyrosyl-L- α -aspartyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 38 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2000:223526 HCAPLUS

DOCUMENT NUMBER: 133:39768

TITLE: Substrate specificity and inhibition studies of human serotonin N-acetyltransferase

Updated Search


09890219

AUTHOR(S): Ferry, Gilles; Loynel, Armelle; Kucharczyk, Nathalie;
Bertin, Sophie; Rodriguez, Marianne; Delagrangé,
Philippe; Galizzi, Jean-Pierre; Jacoby, Edgar;
Volland, Jean-Paul; Lesieur, Daniel; Renard, Pierre;
Canet, Emmanuel; Fauchère, Jean-Luc; Boutin, Jean A.
CORPORATE SOURCE: Division de Pharmacologie Moléculaire et Cellulaire,
Institut de Recherches Servier, Croissy sur Seine,
78290, Fr.
SOURCE: Journal of Biological Chemistry (2000), 275(12),
8794-8805
CODEN: JBCHA3; ISSN: 0021-9258
PUBLISHER: American Society for Biochemistry and Molecular
Biology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Arylalkylamine N-acetyltransferase (AANAT) catalyzes the reaction of serotonin with acetyl-CoA to form N-acetylserotonin and plays a major role in the regulation of the melatonin circadian rhythm in vertebrates. In the present study, the human cloned enzyme has been expressed in bacteria, purified, cleaved, and characterized. The specificity of the human enzyme toward substrates (natural as well as synthetic arylethylamines) and cosubstrates (essentially acyl homologs of acetyl-CoA) has been investigated. Peptide combinatorial libraries of tri-, tetra-, and pentapeptides with various amino acid compns. were also screened as potential sources of inhibitors. We report the findings of several peptides with low micromolar inhibitory potency. For activity measurement as well as for specificity studies, an original and rapid method of anal. was developed. The assay was based on the separation and detection of N-[3H]acetylarylethylamine formed from various arylethylamines and tritiated acetyl-CoA, by means of high performance liquid chromatog. with radio-chemical detection. The assay proved to be robust and flexible, could accommodate the use of numerous synthetic substrates, and was successfully used throughout this study. We also screened a large number of pharmacol. bioamines among which only one, tranlylcypromine, behaved as a substrate. The synthesis and survey of simple arylethylamines also showed that AANAT has a large recognition pattern, including compds. as different as phenyl-, naphthyl-, benzothienyl-, or benzofuranyl-ethylamine derivs. An extensive enzymic study allowed us to pinpoint the amino acid residue of the pentapeptide inhibitor, S 34461, which interacts with the cosubstrate-binding site area, in agreement with an in silico study based on the available coordinates of the hAANAT crystal.

IT 274918-26-2
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(substrate specificity and inhibition studies of human serotonin N-acetyltransferase)
RN 274918-26-2 HCAPLUS
CN L-Tyrosine, 3-benzo[b]thien-3-yl-L-alanyl-4-fluoro-L-phenylalanyl-3-cyclohexyl-L-alanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 70 THERE ARE 70 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 39 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2000:133716 HCAPLUS
 DOCUMENT NUMBER: 132:180873
 TITLE: Preparation of pentapeptide LHRH analogs
 INVENTOR(S): Haviv, Fortuna; Dwight, Wesley; Greer, Jonathan
 PATENT ASSIGNEE(S): Abbott Laboratories, USA
 SOURCE: PCT Int. Appl., 43 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000009545	A1	20000224	WO 1999-US18476	19990812
W: CA, JP, MX				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 6297354	B1	20011002	US 1999-373180	19990812
PRIORITY APPLN. INFO.:			US 1998-132999	A 19980812
			US 1998-96292P	P 19980812
OTHER SOURCE(S):		MARPAT 132:180873		
GI				



AB Pentapeptide LHRH analogs of formula R-A-B-C-D-E-R1 [R = Q; X = H, alkyl, alkoxy, halide; Y = H, alkyl; n = 1-3; A = 3-(1-naphthyl)-D-alanyl(D-1NaI), 3-(1-naphthyl)-L-alanyl, D-Trp, Gly, etc.; B = Ser, Gly; C = NMeTyr, NMePhe, Sar, Arg, Lys(Nε-nicotinoyl), etc.; or B and C together form and amino acid derivative; D = D-Lys(Nε-nicotinyl),

D-Arg, D-Cit, Phe, Gly, etc.; E = cyclohexylalanyl, Gly, Leu, NMeLeu; or D and E together form an amino acid derivative; R1 = NH(CH2)lR2, NR3(CH2)mNHR4, NH(CH2)rNR5R6, NH(CH2)pNHC(:NH)NH2; l = 0-10; m = 1-2; r and p = 1-10; R2 = H, OH, NH2, CONH2, Me, Ph; R3 = H, Me, Et; R4 = H, Me, NH2, CONH2; R5 and R6 taken together with the nitrogen atom to which each is attached form a ring, e.g. pyrrolyl, piperidiny, morpholinyl, pyridyl, etc.], or their pharmaceutically acceptable salts, esters, or prodrugs, were prepared as LHRH antagonists. Thus, 4-F-Phenylpropionyl-D-1Nal-Ser-NMeTyr-D-Lys(Nε-nicotinyl)-Leu-NHCH2CH2-(1-pyrrolidine), prepared by standard solid phase peptide synthesis methods and isolated as the trifluoroacetate salt, antagonized LHRH with pA2 = 9.91. The title compds. are useful in the treatment of disease conditions which are mediated by reproductive hormones, e.g. benign prostate hyperplasia, prostate tumors, breast and ovaries tumors, cryptorchidism, hirsutism, gastric motility disorders, dysmenorrhea, and endometriosis.


IT 259273-45-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of pentapeptide LHRH analogs)

RN 259273-45-5 HCAPLUS

CN L-Tyrosine, N-[3-(4-fluorophenyl)-1-oxopropyl]-O-(phenylmethyl)-L-seryl-O-[(2,6-dichlorophenyl)methyl]-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 40 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1999:718961 HCAPLUS
 DOCUMENT NUMBER: 131:346531
 TITLE: antiangiogenic kringle 5 peptide fragments of plasminogen for therapeutic control of angiogenesis
 INVENTOR(S): Davidson, Donald J.
 PATENT ASSIGNEE(S): Abbott Laboratories, USA
 SOURCE: U.S., 40 pp., Cont.-in-part of U.S. 5,801,146.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

09890219

US 5981484	A	19991109	US 1997-832087	19970403
US 5801146	A	19980901	US 1996-643219	19960503
EP 910571	A2	19990428	EP 1997-925478	19970505
EP 910571	B1	20050720		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI				
CN 1223690	A	19990721	CN 1997-195989	19970505
BR 9708911	A	19990803	BR 1997-8911	19970505
HU 9903530	A2	20000228	HU 1999-3530	19970505
HU 224827	B1	20060228		
US 6057122	A	20000502	US 1997-851350	19970505
NZ 332319	A	20000929	NZ 1997-332319	19970505
JP 2002502235	T	20020122	JP 1997-540162	19970505
AT 299888	T	20050815	AT 1997-925478	19970505
EP 1612272	A2	20060104	EP 2005-106596	19970505
EP 1612272	A3	20070502		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, IE, FI, PL

ES 2246513	T3	20060216	ES 1997-925478	19970505
US 6699838	B1	20040302	US 1997-924287	19970905
US 5972896	A	19991026	US 1998-131995	19980811
US 6251867	B1	20010626	US 1998-132154	19980811
KR 2000010739	A	20000225	KR 1998-708851	19981103
HK 1021191	A1	20060519	HK 1999-104850	19991027
US 2004138127	A1	20040715	US 2004-753646	20040108

PRIORITY APPLN. INFO.:

US 1996-643219	A2	19960503
US 1997-832087	A	19970403
EP 1997-925478	A3	19970505
US 1997-851350	A2	19970505
WO 1997-US7700	W	19970505
US 1997-924287	A1	19970905

AB Mammalian kringle 5 peptide fragments that can inhibit angiogenesis are described for treating angiogenic diseases. Kringle 5 peptide fragments were manufactured either by proteolytic cleavage of plasminogens from various species or synthesized by standard Fmoc chemical. The inhibition of stimulated proliferation and migration by kringle 5 peptide fragments was both potent and specific to the bovine endothelial cells but not normal or tumor cells. Methods and compns. for inhibiting angiogenic diseases are also proposed.

IT 199664-87-4P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)



(synthesis of, antiangiogenic kringle 5 peptide; antiangiogenic kringle 5 peptide fragments of plasminogen for therapeutic control of angiogenesis)

RN 199664-87-4 HCAPLUS

CN L-Tyrosinamide, 1-acetyl-L-prolyl-L-arginyl-L-lysyl-L-leucyl-3-iodo-L-tyrosyl-L- α -aspartyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 41 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1999:670109 HCAPLUS
 DOCUMENT NUMBER: 131:295567
 TITLE: Inhibition of Helicobacter pylori proliferation
 INVENTOR(S): Kaneko, Hiroshi; Mitsuma, Terunori; Yamashita, Koichi;
 Morgan, Barry
 PATENT ASSIGNEE(S): Biomeasure, Inc., USA
 SOURCE: U.S., 19 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5968903	A	19991019	US 1998-74117	19980507
WO 9956769	A2	19991111	WO 1999-US10058	19990506
WO 9956769	A3	20001109		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN,

MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
 TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD,
 RU, TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK,
 ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG,
 CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 9939754 A 19991123 AU 1999-39754 19990506

EP 1075273 A2 20010214 EP 1999-922851 19990506

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI

JP 2002513769 T 20020514 JP 2000-546793 19990506

NO 2000005588 A 20010105 NO 2000-5588 20001106

PRIORITY APPLN. INFO.: US 1998-74117 A1 19980507

WO 1999-US10058 W 19990506

OTHER SOURCE(S): MARPAT 131:295567

AB The present invention is directed to a method of using somatostatin or a somatostatin agonist to inhibit the proliferation of *Helicobacter pylori* (*H. pylori*), which comprises administering to a patient in need thereof an effective amount of said somatostatin or somatostatin agonist. Preferably, a somatostatin sub-type receptor 2 (SSTR-2) selective somatostatin agonist is administered in a method of this invention. The inhibition of *H. pylori* proliferation is useful in treating various gastroduodenal diseases such as peptic ulcers, gastric cancer and gastric lymphoma.

IT 113294-83-0

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(inhibition of *Helicobacter pylori* proliferation with somatostatin or a somatostatin agonist)


RN 113294-83-0 HCAPLUS

CN L-Threoninamide, 2,3,4,5,6-pentafluoro-D-phenylalanyl-L-cysteinyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-cysteinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





REFERENCE COUNT: 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 42 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:578704 HCAPLUS

DOCUMENT NUMBER: 132:102593

TITLE: Agonist-antagonist structure-activity relationships of thrombin receptor tethered ligand peptide

AUTHOR(S): Fujita, T.; Nose, T.; Nakajima, M.; Inoue, Y.; Nakamura, N.; Inoue, T.; Costa, T.; Shimohigashi, Y.

CORPORATE SOURCE: Laboratory of Biochemistry, Department of Chemistry, Faculty of Science, Kyushu University, Fukuoka, 812-8581, Japan

SOURCE: Peptide Science: Present and Future, Proceedings of the International Peptide Symposium, 1st, Kyoto, Nov. 30-Dec. 5, 1997 (1999), Meeting Date 1997, 202-204. Editor(s): Shimonishi, Yasutsugu. Kluwer: Dordrecht, Neth.

CODEN: 68BYA5

DOCUMENT TYPE: Conference

LANGUAGE: English

AB In order to obtain an effective antagonist of thrombin receptor, we have designed several SFLLRNP analogs that could be expected to establish new interaction with the receptor.

IT 255837-51-5

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
(agonist-antagonist structure-activity relationships of thrombin receptor tethered ligand peptide)


RN 255837-51-5 HCAPLUS

CN L-Tyrosinamide, 4-fluoro-N-(3-mercapto-1-oxopropyl)-L-phenylalanyl-3-

09890219

cyclohexyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 43 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:453428 HCAPLUS

DOCUMENT NUMBER: 131:267135

TITLE: Radioligand binding assay of cholecystokinin receptor in rat cerebral cortex

AUTHOR(S): Xiang, Peng; Chen, Manling; Tan, Tianzhi; Shi, Yuhong
CORPORATE SOURCE: School of Basic Medical Sciences, WCUMS, Chengdu, 610041, Peop. Rep. China

SOURCE: Huaxi Yike Daxue Xuebao (1999), 30(2), 214-216
CODEN: HYDXET; ISSN: 0257-7712

PUBLISHER: Huaxi Yike Daxue

DOCUMENT TYPE: Journal

LANGUAGE: Chinese

AB A radioligand binding assay system for determining the characterization of CCK receptor is presented. Using Bolton-Hunter reagent, the authors prepared a biol. active, specific ^{125}I -BH-CCK8. The iodination mixture was then transferred to a column of Sephadex G-25 and examined by silica TLC. Its specific activity and radiochem. purity were 3.4 TBq/mmol and 96%, resp. Binding of ^{125}I -BH-CCK8 to the membrane of rat cerebral cortex was rapid, reversible, time-temperature dependent, saturable and specific. The labeled

CCK was shown to have biol. activity as measured by the CCK receptor radioassay. Under the authors' laboratory conditions, the CCK binding required 1 h to reach equilibrium at 4°. The authors chose polyethylene glycol 6000 and γ -globulin protein for the separation of B and F. Scatchard plot of CCK binding was linear with a K_d value of 1.098 nmol/L and B_{max} of 197.5 fmol/mg protein. The results of this study support the view that CCK may function as a regulatory peptide in brain and hence may be of use for clarifying the CCK receptor's function in central nervous system.

IT 79672-09-6

RL: ANT (Analyte); PRP (Properties); ANST (Analytical study)
(radioligand binding assay of cholecystokinin receptor in rat cerebral cortex)

RN 79672-09-6 HCAPLUS

CN Cholecystokinin-8 (swine), N-[3-[4-hydroxy-3-(iodo- ^{125}I)phenyl]-1-


Updated Search

09890219


oxopropyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L11 ANSWER 44 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

Updated Search

09890219



ACCESSION NUMBER: 1999:383879 HCAPLUS
DOCUMENT NUMBER: 131:175140
TITLE: Determination of ABT-861 by high-performance liquid chromatography and a model for ion-pair formation with trifluoroacetic acid
AUTHOR(S): Simonzadeh, N.; Levison, B.; Buko, A.; Matuszak, K.; Hanscom, M.
CORPORATE SOURCE: Abbott Laboratories, Abbott Park, IL, 60064-3500, USA
SOURCE: Journal of Chromatographic Science (1999), 37(6), 185-190
CODEN: JCHSBZ; ISSN: 0021-9665
PUBLISHER: Preston Publications
DOCUMENT TYPE: Journal
LANGUAGE: English

AB ABT-861 is a gonadotropin releasing hormone (GnRH) antagonist candidate drug synthesized at Abbott Labs. for use in medical conditions responsive to hormonal manipulation (e.g., prostate cancer in elderly males, endometriosis in females, and central precocious puberty in children). A HPLC method employing gradient elution with UV detection is developed for the assay of ABT-861 and determination of impurities in bulk powder and injectable formulations. The chromatog. conditions employed included the use of a 250 + 4.6 mm, 5- μ m ODS Vydac HPLC column at 35°, an MeCN-H₂O (0.1% TFA in each phase) eluent, and a 60-min run time using UV detection. The chromatog. conditions are used for the determination of ABT-861 and its degradation products and manufacturing impurities in the bulk powder and injectable formulations. The limit of detection was approx. 9 ng at 225 nm. Method validation includes linearity of detector response with amount injected, precision, and standard addition-recovery data. Under the chromatog. conditions employed, diastereomeric and manufacturing impurities and degradation products are separated from ABT-861, demonstrating that the method is stability-indicating. Thus, the current method is suitable for the routine anal. of ABT-861 and related impurities, providing good selectivity and sensitivity. (c) 1999 Preston Publications.

IT 238080-44-9
RL: ANT (Analyte); FMU (Formation, unclassified); ANST (Analytical study); FORM (Formation, nonpreparative)
(determination of ABT-861 by HPLC and model for ion-pair formation with trifluoroacetic acid)

RN 238080-44-9 HCAPLUS
CN D-Alaninamide, N-[[[(2S)-tetrahydro-2-furanyl]carbonyl]glycyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-L-seryl-N-methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-methylethyl)-L-lysyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 45 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:353265 HCAPLUS

DOCUMENT NUMBER: 131:166695

TITLE: Structural essentials for agonist-antagonist actions of thrombin receptor tethered-ligand

AUTHOR(S): Nose, Takeru; Fujita, Tsugumi; Morita, Yuki; Costa, Tommaso; Shimohigashi, Yasuyuki

CORPORATE SOURCE: Department of Chemistry, Faculty of Science, Kyushu University, Fukuoka, 812-8581, Japan

SOURCE: Peptide Science (1999), Volume Date 1998, 35th, 217-220

CODEN: PSCIFQ; ISSN: 1344-7661


PUBLISHER: Protein Research Foundation

DOCUMENT TYPE: Journal
LANGUAGE: English

IT 238756-24-6

RN 238756-24-6 HCAPLUS

Absolute stereochemistry.
Double bond geometry as shown.



L11 ANSWER 46 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:329683 HCAPLUS

DOCUMENT NUMBER: 131:115001

TITLE: Recognition of an MHC class I-restricted antigenic peptide can be modulated by para-substitution of its buried tyrosine residues in a TCR-specific manner

AUTHOR(S): Saito, Naoyuki G.; Chang, Hsiu-Ching; Paterson, Yvonne
CORPORATE SOURCE: Department of Microbiology and Eldridge Reeves Johnson
Foundation for Molecular Biophysics, University of
Pennsylvania School of Medicine, Philadelphia, PA,
19104, USA

SOURCE: Journal of Immunology (1999), 162(10), 5998-6008

CODEN: JOIMA3; ISSN: 0022-1767

PUBLISHER: American Association of Immunologists

DOCUMENT TYPE: Journal


LANGUAGE: English



AB Conformational dependence of TCR contact residues of the H-2Kb mol. on the two buried tyrosine side chains of the vesicular stomatitis virus (VSV)-8 peptide was investigated by systematic substitutions of the tyrosines with

Updated Search

IT	232616-70-5D, complexes with MHC class I RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study) (cytotoxic T-cell recognition of MHC class I/peptide complexes is modulated by complex contact surface for TCR as altered by anchor residue structure in binding pocket)
RN	232616-70-5 HCAPLUS
CN	L-Leucine, L-arginylglycyl-4-fluoro-L-phenylalanyl-L-valyl-L-tyrosyl-L-glutaminylglycyl- (9CI) (CA INDEX NAME)

PAGE 1-A





REFERENCE COUNT: 86 THERE ARE 86 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 47 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1998:344369 HCAPLUS
 DOCUMENT NUMBER: 129:16399
 TITLE: Conformationally constrained LH-RH analogs, their uses and pharmaceutical compositions containing them
 INVENTOR(S): Delansorne, Remi; Paris, Jacques
 PATENT ASSIGNEE(S): Laboratoire Theramex S.A., Monaco
 SOURCE: Eur. Pat. Appl., 32 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 842946	A1	19980520	EP 1996-402441	19961114
R: FR				
CA 2270158	A1	19980522	CA 1997-2270158	19971112
WO 9821229	A1	19980522	WO 1997-EP6322	19971112
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AU 9854817	A	19980603	AU 1998-54817	19971112
EP 937101	A1	19990825	EP 1997-951183	19971112
EP 937101	B1	20001011		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
CN 1237979	A	19991208	CN 1997-199752	19971112
BR 9713060	A	20000411	BR 1997-13060	19971112
HU 200000249	A2	20000728	HU 2000-249	19971112
HU 200000249	A3	20001128		
AT 196914	T	20001015	AT 1997-951183	19971112
NZ 335454	A	20001027	NZ 1997-335454	19971112
JP 2001503444	T	20010313	JP 1998-522187	19971112
ZA 9710254	A	19980528	ZA 1997-10254	19971113
NO 9902298	A	19990708	NO 1999-2298	19990512
MX 9904512	A	20000531	MX 1999-4512	19990514

09890219

US 6153587	A	20001128	US 1999-317125	19990524
PRIORITY APPLN. INFO.:			EP 1996-402441	A 19961114
			WO 1997-EP6322	W 19971112

OTHER SOURCE(S): MARPAT 129:16399

AB LH-RH peptide analogs V-W-X-SPL-Y-Pro-Z [V is the peptide A1A2, where A1 is pGlu, AcSar, or an aromatic D-amino acid and A2 is a direct bond, His, D-Phe, D-pFPhe, or D-pClPhe; W is an aromatic L- or D-amino acid; X is the dipeptide A3A4, where A3 is Ala, Thr, Ser, D-Ser, Ser(OBzl), or MeSer and A4 = Tyr, Phe, cis-3-(4-pyrazinylcarbonylaminocyclohexyl)alanine, L- or D-Ne-picolinoylllysine, -Ne-nicotinoylllysine, or -Ne-isopropoylllysine; SPL is a spirolactam; Y is the dipeptide A5A6, where A5 is an amino acid with an alkyl or cycloalkyl side chain and A6 is (un)substituted L- or D-Arg, -homoarginine, -Lys, -homolysine, -Orn, -citrulline, -homocitrulline, or p-aminophenylalanine; Z is GlyNH₂, D-Ala-NH₂, azaglycinamide, (un)substituted alkylamino] were prepared which have excellent affinity for LH-RH receptors. Thus, pGlu-His-Trp-Ser-Tyr-SPL-Leu-Arg-Pro-NHEt, prepared by the solid-phase method, showed affinity (pK_i = 8.94) for the LH-RH receptor.

IT 207607-70-3P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(conformationally constrained LH-RH analogs)


RN 207607-70-3 HCAPLUS

CN Glycinamide, 5-oxo-L-prolyl-L-histidyl-4-chloro-L-phenylalanyl-L-seryl-L-tyrosyl-(α S,5S)- α -(2-methylpropyl)-6-oxo-1,7-diazaspiro[4.4]nonane-7-acetyl-L-arginyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 48 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1998:163467 HCAPLUS

DOCUMENT NUMBER: 128:226683

TITLE: Method of inhibiting fibrosis with a somatostatin agonist

INVENTOR(S): Culler, Michael D.; Kasprzyk, Philip G.

PATENT ASSIGNEE(S): Biomeasure Incorporated, USA; Culler, Michael D.; Kasprzyk, Philip G.

SOURCE: PCT Int. Appl., 61 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9808529	A1	19980305	WO 1997-US14154	19970827
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2264309	A1	19980305	CA 1997-2264309	19970827
AU 9741490	A	19980319	AU 1997-41490	19970827
AU 726731	B2	20001116		
EP 938328	A1	19990901	EP 1997-939392	19970827
EP 938328	B1	20060412		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
CN 1229357	A	19990922	CN 1997-197671	19970827

09890219

HU 9903918	A2	20000428	HU 1999-3918	19970827
JP 2001500483	T	20010116	JP 1998-511678	19970827
EP 1574219	A2	20050914	EP 2005-76124	19970827
EP 1574219	A3	20060426		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI

AT 322905	T	20060415	AT 1997-939392	19970827
ES 2258282	T3	20060816	ES 1997-939392	19970827
ZA 9707783	A	19990301	ZA 1997-7783	19970829
US 6268342	B1	20010731	US 1999-254097	19990510
US 2005222025	A1	20051006	US 2004-935593	20040907

PRIORITY APPLN. INFO.:

US 1996-705790	A2	19960830
EP 1997-939392	A3	19970827
WO 1997-US14154	W	19970827
US 1999-254097	A3	19990510
US 2001-761605	A3	20010116

OTHER SOURCE(S): MARPAT 128:226683

AB The present invention relates to a method of inhibiting fibrosis in a patient. The method comprises administering a therapeutically effective amount of a somatostatin, a somatostatin agonist or a pharmaceutically acceptable salt thereof to said patient.

IT 113294-83-0

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(method of inhibiting fibrosis with a somatostatin agonist)

RN 113294-83-0 HCAPLUS


CN L-Threoninamide, 2,3,4,5,6-pentafluoro-D-phenylalanyl-L-cysteinyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-cysteinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



Updated Search



REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 49 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1997:740418 HCAPLUS

DOCUMENT NUMBER: 128:43873

TITLE: Antiangiogenic peptides, polypeptides containing them, and methods for inhibiting angiogenesis

INVENTOR(S): Davidson, Donald J.; Wang, Jieyi; Gubbins, Earl J.

PATENT ASSIGNEE(S): Abbott Laboratories, USA

SOURCE: PCT Int. Appl., 78 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9741824	A2	19971113	WO 1997-US7700	19970505
WO 9741824	A3	19980108		
W: AU, BR, CA, CN, CZ, HU, IL, JP, KR, MX, NZ				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5801146	A	19980901	US 1996-643219	19960503
CA 2253243	A1	19971113	CA 1997-2253243	19970505
AU 9730606	A	19971126	AU 1997-30606	19970505
AU 724077	B2	20000914		
EP 910571	A2	19990428	EP 1997-925478	19970505
EP 910571	B1	20050720		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI				
CN 1223690	A	19990721	CN 1997-195989	19970505
BR 9708911	A	19990803	BR 1997-8911	19970505

09890219

HU 9903530	A2	20000228	HU 1999-3530	19970505
HU 224827	B1	20060228		
NZ 332319	A	20000929	NZ 1997-332319	19970505
JP 2002502235	T	20020122	JP 1997-540162	19970505
AT 299888	T	20050815	AT 1997-925478	19970505
HK 1021191	A1	20060519	HK 1999-104850	19991027

PRIORITY APPLN. INFO.:

US 1996-643219	A	19960503
US 1997-832087	A	19970403
WO 1997-US7700	W	19970505

AB Mammalian kringle 5 fragments and kringle 5 fusion proteins are disclosed as compds. for treating angiogenic diseases. Methods and compns. for inhibiting angiogenic diseases are also disclosed.

IT 199664-87-4P

RL: PNU (Preparation, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(antiangiogenic peptides, polypeptides containing them, and methods for inhibiting angiogenesis)

RN 199664-87-4 HCAPLUS


CN L-Tyrosinamide, 1-acetyl-L-prolyl-L-arginyl-L-lysyl-L-leucyl-3-iodo-L-tyrosyl-L- α -aspartyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B




L11 ANSWER 50 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

Updated Search

09890219

ACCESSION NUMBER: 1997:547277 HCAPLUS
 DOCUMENT NUMBER: 127:162122
 TITLE: Preparation of 5-amino-4-hydroxyhexanoic acid derivatives for treatment of AIDS
 INVENTOR(S): Bold, Guido; Lang, Marc; Fassler, Alexander; Capraro, Hans-georg; Bhagwat, Shripad; Schneider, Peter; Hoogevest, Peter van
 PATENT ASSIGNEE(S): Ciba-Geigy Corp., USA
 SOURCE: U.S., 98 pp., Cont.-in-part of U.S. Ser. No. 941,595, abandoned.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5643878	A	19970701	US 1994-207646	19940308
ZA 9206938	A	19940311	ZA 1992-6938	19920911
CN 1089269	A	19940713	CN 1993-100044	19930104
PRIORITY APPLN. INFO.:			CH 1991-2689	A 19910912
			CH 1992-890	A 19920327
			CH 1992-2007	A 19920625
			US 1992-941595	B2 19920908
			CH 1992-772	A 19930311
OTHER SOURCE(S):		MARPAT 127:162122		
GI				



I

AB Peptides I [A1, B1 = bond, amino acid residue; A2 = amino acid residue; R1 = H, alkoxycarbonyl, or (un)substituted benzyloxycarbonyl; R2, R3 = (un)substituted Ph or cyclohexyl; R4R5N = (un)substituted morpholino] were prepared for the treatment of AIDS. Thus, 5(S)-Boc-amino-4(S)-hydroxy-6-cyclohexyl-2(R)-(p-fluorophenylmethyl)hexanoyl-L-Val-L-Phe-morpholin-4-ylamide (Boc = tert-butoxycarbonyl) was prepared via peptide coupling in solution


IT 150609-45-3P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of aminohydroxyhexanoic acid derivs. for treatment of AIDS)

RN 150609-45-3 HCAPLUS

CN Carbamic acid, [1-(cyclohexylmethyl)-2-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-4-[(4-fluorophenyl)methyl]-5-[[1-[[[1-[(4-methoxyphenyl)methyl]-2-(4-morpholinyl)-2-oxoethyl]amino]carbonyl]-2-methylpropyl]amino]-5-oxopentyl]-, 1,1-dimethylethyl ester, [1S-[1R*,2R*,4S*,5[R*(R*)]]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



L11 ANSWER 51 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1996:721028 HCAPLUS

DOCUMENT NUMBER: 126:55003

TITLE: Radiolabeled ligands specific for the G protein-coupled state of neurotensin receptors

AUTHOR(S): Gaudriault, Georges; Zsuzsanna, Nicole; Vincent, Jean-Pierre

CORPORATE SOURCE: Institut Pharmacologie Moléculaire Cellulaire, Université Nice Sophia Antipolis, Valbonne, Fr.

SOURCE: Journal of Neurochemistry (1996), 67(6), 2590-2598
CODEN: JONRA9; ISSN: 0022-3042

PUBLISHER: Lippincott-Raven

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Radiolabeled analogs of neuromedin N have been prepared by acylation of the α , ϵ 1, and ϵ 2 amino groups of [Lys2]neuromedin N (Lys-Lys-Pro-Tyr-Ile-Leu) either with the 125I-labeled Bolton-Hunter reagent or with N-succinimidyl[2,3-³H]propionate. The binding properties of the purified analogs toward newborn mouse brain homogenate or toward membranes of cells transiently (COS) or permanently (AA1) transfected with the cloned rat brain neurotensin receptor cDNA were evaluated and compared with those of radiolabeled neurotensin. The α -modified analog of [Lys2]neuromedin N behaves exactly like neurotensin in these binding expts., whereas the ϵ 1- and ϵ 2-modified analogs selectively recognize the fraction of neurotensin binding sites that is sensitive to GTP γ S. The proportion of neurotensin receptors coupled to GTP binding proteins is approx.50% in membranes of newborn mouse brain or of AA1 cells that respond to neurotensin by an increase of the intracellular inositol trisphosphate concentration. By contrast, membranes of transiently transfected COS cells that do not respond to neurotensin exhibit very low levels of GTP-sensitive receptors labeled with the ϵ 1- or ϵ 2-modified analogs. These radiolabeled peptides offer new tools to selectively detect active neurotensin receptors.

IT 185154-30-7

09890219


RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(radiolabeled ligands specific for G protein-coupled state of neurotensin receptors)

RN 185154-30-7 HCAPLUS


2-13-Neurotensin (cattle), N-[3-[4-hydroxy-3-(iodo-125I)phenyl]-1-oxopropyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 2-A



Updated Search

L11 ANSWER 52 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1996:175605 HCAPLUS

DOCUMENT NUMBER: 124:233153

TITLE: Preparation of compound bearing two
2,6-diiodophenol-4-yl groups and diagnostic drug for
iodine allergy

INVENTOR(S): Sugihara, Yoshiki; Shionoya, Hiroshi; Yamatsu, Kiyomi

PATENT ASSIGNEE(S): Eisai Co., Ltd., Japan

SOURCE: PCT Int. Appl., 38 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese



FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9532173	A1	19951130	WO 1995-JP997	19950524
W: CA, JP, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
JP 3349513	B2	20021125	JP 1995-530201	19950524
US 5780509	A	19980714	US 1997-737625	19970129
PRIORITY APPLN. INFO.:			JP 1994-109374	A 19940524
			WO 1995-JP997	W 19950524

OTHER SOURCE(S): MARPAT 124:233153

GI



AB A compound bearing at least two 2,6-diiodophenol-4-yl (antigen determinant) groups (I; X = monovalent atom or group; Y = bivalent atom or group), particularly a peptide composed of 2 to 8 iodinated tyrosine mols. condensed with each other (II; X, X1 = monovalent atom or group; R = H, iodo; n = 0-6; wherein R is selected such that the peptide contains more

than two 2,6-diiodophenol-4-yl group), is prepared This compound is useful as a diagnostic drug for iodine allergy, particularly against a x-ray imaging contrast agent for human urethra or blood vessels. The diagnostic method, which is based on the finding that the 2,6-diiodophenol-4-yl group is an antigenic determinant for iodine allergy, comprises administering the drug II (0.5-50 µg) to the patient to examine the intracutaneous reaction. Thus, 250 mg H-Tyr-Tyr-Tyr-OH was added to 50 mM (NH₄)₂CO₃ buffer solution (pH 9.4, 20 mL) and dissolved by adding 1 N NaOH, followed by adding portion wise 0.5 M KI (5.6 mL) and 1 N NaOH, and the resulting mixture was stirred at room temperature for 10 min, treated with sodium thiosulfate to decompose unreacted iodine, made pH 3 with 1 N HCl, and extracted with EtOAc to give 236 mg hexaiodotritityrosine II (R = iodo, X = H₂N, X₁ = CO₂H, n = 1). When the latter compound (0.1 mL of 1 µg/mL solution) was s.c. administered to iodine-sensitized guinea pigs, active cutaneous anaphylaxis was induced.

IT 174608-41-4P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of peptide containing diiodotyrosine as diagnostic agents for iodine allergy)

RN 174608-41-4 HCAPLUS

CN L-Tyrosine, 3,5-diiodo-L-tyrosyl-3,5-diiodo-L-tyrosyl-3,5-diiodo- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 53 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1995:958268 HCAPLUS

DOCUMENT NUMBER: 123:350253

TITLE: Aerosol drug formulations containing vitamin E

INVENTOR(S): Fu, Lu Mou-ying; Gupta, Pramod K.; Adjei, Akwete L.

PATENT ASSIGNEE(S): Abbott Laboratories, USA

SOURCE: PCT Int. Appl., 18 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.

KIND

DATE

APPLICATION NO.

DATE

WO 9524892	A1	19950921	WO 1995-US2764	19950302
W: AU, CA, JP, KR, MX				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2183557	A1	19950921	CA 1995-2183557	19950302
AU 9519804	A	19951003	AU 1995-19804	19950302
AU 709783	B2	19990909		
JP 09510445	T	19971021	JP 1995-524061	19950302
EP 804157	A1	19971105	EP 1995-912746	19950302
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE				
PRIORITY APPLN. INFO.:			US 1994-212472	A 19940314
			WO 1995-US2764	W 19950302

AB Pharmaceutical compns. for aerosol delivery are disclosed comprising (a) a medicament, (b) a non-chlorofluorocarbon propellant, and (c) tocopherol or a pharmaceutically acceptable derivative thereof, as well as a method for preparing such compns. in which unwanted aggregation of the medicament is prevented without the use of surfactants or cosolvents. Pharmaceutical aerosols containing leuprolide acetate in 0.1% d- α tocopheryl acetate (I) and 10mL HFC-134a were prepared having good dispersion quality as compared with controls without I which had poor dispersion quality.

IT 170929-31-4


RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(aerosol drug formulations containing vitamin E)


RN 170929-31-4 HCAPLUS

CN D-Alaninamide, N-acetyl-3-(1-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-O-(1-oxohexadecyl)-D-seryl-N-methyl-L-tyrosyl-N6-(3-pyridinylcarbonyl)-D-lysyl-L-leucyl-N6-(1-methylethyl)-L-lysyl-L-prolyl-(9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A






PATENT INFORMATION:

GI



I



II

AB Title compds. [I; T = R6CO; R6 = (substituted) hydrocarbyl in which ≥ 1 C atom is replaced by a heteroatom; R1 = H, alkoxycarbonyl, heterocyclylcarbonyl, (substituted) benzyloxycarbonyl, heterocyclyloxycarbonyl, etc.; A1, B1 = bond, amino acid residue; R2, R3 = (substituted) Ph, cyclohexyl; A2 = amino acid residue; A1A2 = dipeptide residue whose central amide bond is reduced; NR4R5 = (substituted) morpholino, thiomorpholino], were prepared Title compound II was prepared by solution phase coupling reactions. I inhibited HIV-1 protease with $IC_{50} = 10^{-7}$ - 10^{-9} M.


IT 150608-33-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of dipeptide derivs. of 5-amino-4-hydroxyhexanoic acid as HIV protease inhibitors)

RN 150608-33-6 HCAPLUS

CN Carbamic acid, [4-[(4-fluorophenyl)methyl]-2-hydroxy-5-[[1-[[[1-[(4-methoxyphenyl)methyl]-2-(4-morpholinyl)-2-oxoethyl]amino]carbonyl]-2-methylpropyl]amino]-5-oxo-1-(phenylmethyl)pentyl]-, 1,1-dimethylethyl ester, [1S-[1R*,2R*,4S*,5[R*(R*)]]]- (9CI) (CA INDEX NAME)




L11 ANSWER 55 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1995:666961 HCAPLUS
 DOCUMENT NUMBER: 123:48053
 TITLE: Reduced-Size Antagonists of Luteinizing
 Hormone-Releasing Hormone Active in Vitro
 AUTHOR(S): Janecka, Anna; Janecki, Tomasz; Bowers, Cyril;
 Folkers, Karl
 CORPORATE SOURCE: Institute for Biomedical Research, University of
 Texas, Austin, TX, 78705, USA
 SOURCE: Journal of Medicinal Chemistry (1995), 38(15), 2922-4
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English



AB A series of reduced-size analogs of LHRH was designed with the length
 varying from nine to two amino acids. These compds. were tested in vitro
 for the LH suppression in cultured rat pituitary cells treated with 1 ng
 of LHRH. The best analogs were also tested in vivo for their
 antioviulatory activity in rats. It appeared that terminal amino acids as
 well as the presence of Arg or Ilys (Nε-isopropyllysine) in the
 sequence are both crucial for the antagonism. The most potent antagonist
 in this series was a heptapeptide, Ac-D-Nal-Ser-Tyr-D-Nal-Leu-Arg-ProNH₂,
 which completely inhibited LH release at the concentration 0.1 μg/mL and
 inhibited ovulation at 1000 μg/rat. For fragments shorter than
 heptapeptide the inhibition of LH release was observed at 100 μg/mL
 concentration
 of the analog.

IT 162152-59-2
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); PRP (Properties); BIOL (Biological study)
 (reduced-size antagonists of LH-releasing hormone active in vitro)
 RN 162152-59-2 HCAPLUS
 CN D-Alaninamide, N-acetyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-
 phenylalanyl-L-seryl-L-tyrosyl-3-(2-naphthalenyl)-D-alanyl-L-leucyl-L-
 arginyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





L11 ANSWER 56 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1995:568949 HCAPLUS
DOCUMENT NUMBER: 123:1202
TITLE: Three-Dimensional Quantitative Structure-Activity Relationships of Somatostatin Analogs. 1. Comparative Molecular Field Analysis of Growth Hormone Release-Inhibiting Potencies
AUTHOR(S): Hocart, Simon J.; Reddy, Vik; Murphy, William A.; Coy, David H.
CORPORATE SOURCE: School of Medicine, Tulane University, New Orleans, LA, 70112, USA
SOURCE: Journal of Medicinal Chemistry (1995), 38(11), 1974-89
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The previous work on the structure-activity relation of somatostatin and that of many others has generated a large database of analogs with different biol. activities and receptor affinities. This present work is an investigation of the growth hormone release-inhibiting potencies of somatostatin analogs by the 3-dimensional quant. structure-activity paradigm, comparative mol. field anal. (CoMFA). A total of 64 analogs were modeled in SYBYL using structural information from 2 NMR studies. The mols. were aligned by a root-mean-square fit of atoms and field-fit of the steric and electrostatic mol. fields and the resulting databases

analyzed by partial least squares anal. with cross-validation to extract the optimum number of components. The anal. was then repeated without cross-validation to give the final QSAR models. Preliminary investigations with the CoMFA models led to the synthesis of a new somatostatin analog. This compound together with 5 other newly synthesized compds. not included in the original training sets were used to test the predictive ability of the CoMFA models. Two models with good predictive powers are presented.

IT 150155-65-0, BIM 23067


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(three-dimensional QSARs of somatostatin analogs and comparative mol. field anal. of growth hormone release-inhibiting potencies)


RN 150155-65-0 HCAPLUS

CN D-Phenylalaninamide, 4-chloro-D-phenylalanyl-L-alanyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A





L11 ANSWER 57 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1995:480169 HCAPLUS
 DOCUMENT NUMBER: 122:240447
 TITLE: Preparation of peptideamide analogs as tachykinin antagonists.
 INVENTOR(S): Pieper, Helmut; Austel, Volkhard; Jung, Birgit; Buerger, Erich; Entzeroth, Michael
 PATENT ASSIGNEE(S): Karl Thomas GmbH, Germany
 SOURCE: Ger. Offen., 101 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 4243858	A1	19940630	DE 1992-4243858	19921223
PRIORITY APPLN. INFO.:			DE 1992-4243858	19921223
OTHER SOURCE(S):		MARPAT 122:240447		
GI				



I

AB R4R5NACONHCHR3CXNR1R2 [A = 1,2-cyclopentylene, CHR6; R6 = H, (substituted) alkyl, Ph; R1 = H, (Ph- or pyridyl-substituted) alkyl; R2 = H, (amino- or guanidino-substituted) Ph, pyridyl, (cyclohexyl-, Ph-, or pyridyl-substituted) alkyl, etc.; R1R2N = (substituted) piperazinyl; R3 = H, (phenyl)alkyl, guanidino- or amino-substituted alkyl, aminocarbonylalkyl, etc.; R4 = H, (phenyl)alkyl; R5 = protecting group, (substituted) alkyl, alkanoyl, alkoxycarbonyl, alkylaminocarbonyl, PhCO, naphthylcarbonyl, biphenylcarbonyl, PhSO2, etc.; X = (H, H), O, S; the C atom bearing the R3 substituent is L; the C atom bearing the R6 substituent is D or L], were prepared Thus, title compound I (prepared by solution

phase methods) showed IC50 = 2 nM for neurokinin-1 receptor binding with IM-9 cells. Tablets were prepared containing I.


IT 162177-03-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of, as tachykinin antagonist)

RN 162177-03-9 HCAPLUS

CN L-Lysinamide, 4-amino-3,5-dibromo-N-(1-oxo-4-phenylbutyl)-D-phenylalanyl-N-[2-(2-methoxyphenyl)ethyl]-N6-[(phenylmethoxy)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 58 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1995:408550 HCAPLUS

DOCUMENT NUMBER: 122:161391

TITLE: Preparation of fluorine containing atrial natriuretic peptides

INVENTOR(S): Rakhit, Sumanas; Goghari, Mahesh H.

PATENT ASSIGNEE(S): Bio-Mega/Boehringer Ingelheim Research Inc., Can.

SOURCE: U.S., 19 pp. Cont.-in-part of U.S. 5,095,004.

CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

09890219

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5376635	A	19941227	US 1991-781590	19911023
US 5095004	A	19920310	US 1988-166526	19880314
PRIORITY APPLN. INFO.:			CA 1987-532982	A 19870325
			CA 1987-542192	A 19870715
			US 1988-166526	A2 19880314

OTHER SOURCE(S): MARPAT 122:161391
GI



AB Disclosed herein are derivs. of atrial natriuretic peptides which are characterized by having (at positions 106 and/or 124) a phenylalanyl residue bearing a F or CF3 substituent on the aromatic portion thereof. Said peptides are represented by a general formula [I; R1, R4 = Phe, 2FPhe, 3FPhe, 4FPhe, 2CF3Phe, 3CF3Phe, 4CF3Phe; R2 = Gly, Ala, D-Ala; R3 = Ile, Met; R5 = Tyr, absent; Y = S-(lower alkylene)-CO, R6-Cys; wherein R6 = peptide residue, e.g., H-Ser-Ser and H-Arg-Ser-Ser; provided that when R1 = Phe, R4 ≠ Phe]. Optionally, the exocyclic N-terminal peptide segment and the first cysteinyl residue (at position 105) are replaced by an optionally substituted thioalkanoyl residue. The derivs. possess useful diuretic, natriuretic and antihypertensive activities. Thus, (4FPhe106)hANP(103-125) (II) was prepared by the solid phase method using Boc-Arg(Tos)-PAB-benzhydrylamine resin [PAB = α-(phenylacetamido)benzyl]. In a diuretic assay using normotensive rats, II at 0.5 µg/kg per min urine excretion from 0.42 mL/10 min (control) to 1.27 mL/10 min.

IT 120728-19-0P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of fluorine-containing atrial natriuretic peptides as diuretics, natriuretics, and antihypertensives)
RN 120728-19-0 HCAPLUS
CN Atrial natriuretic peptide-28 (human), 26-(4-fluoro-L-phenylalanine)-(9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

Updated Search

09890219

L11 ANSWER 59 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1995:252000 HCAPLUS

DOCUMENT NUMBER: 122:46704

TITLE: Subtype selectivity of peptide analogs for all five cloned human somatostatin receptors (hsstr 1-5)

AUTHOR(S): Patel, Yogesh C.; Srikant, Coimbatore B.

CORPORATE SOURCE: Fraser Lab., McGill Univ., Montreal, QC, H3A 1A1, Can.

SOURCE: Endocrinology (1994), 135(6), 2814-17

CODEN: ENDOAO; ISSN: 0013-7227

PUBLISHER: Endocrine Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Recent reports (Raynor et al) have claimed the identification of potent somatostatin (SST) agonists exhibiting binding affinities of 1-2 pM and up to 30,000-fold binding selectivity for several of the 5 cloned sstr subtypes. These conclusions, however, are based on binding comparisons of sstr subtypes from different species expressed in different cell lines and studied with different radioligands. To eliminate the effect of species and/or methodol. variations, we have investigated agonist selectivity of 32 synthetic SST analogs for all 5 hsstrs stably expressed in CHO-K1 cells under identical binding conditions. We show that hsstr2, 3, 5 react potently with hexapeptide as well as cyclic and linear octapeptide analogs and belong to a similar sstr subclass. Hsstr1 and 4 react poorly with these analogs and belong to a sep. subclass. The present generation of SST analogs exhibit a modest .apprx. 50-fold increase in binding potency compared to SST-14 for 2 subtypes (hsstr2, 3), and relative selectivity for only 2 subtype (hsstr2) which is at best only 35- fold. The potency and degree of selectivity of these analogs is several orders of magnitude less than that reported earlier and suggests the need for caution in using these compds. as putative superagonists or subtype selective compds. for any of the individual sstrs.



IT 150155-65-0, BIM 23067

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(somatostatin receptor subtype selectivity of)

RN 150155-65-0 HCAPLUS

CN D-Phenylalaninamide, 4-chloro-D-phenylalanyl-L-alanyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 60 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1994:436149 HCAPLUS
DOCUMENT NUMBER: 121:36149
TITLE: In vitro and in vivo Activities of Reduced-Size
Antagonists of Luteinizing Hormone-Releasing Hormone
AUTHOR(S): Haviv, Fortuna; Fitzpatrick, Timothy D.; Nichols,

Charles J.; Bush, Eugene N.; Diaz, Gilbert; Bammert, Gary; Nguyen, A. T.; Johnson, Edwin S.; Knittle, Judith; Greer, Jonathan

CORPORATE SOURCE: TAP Pharmaceutical Products Inc, Abbott Park, IL, 60064, USA

SOURCE: Journal of Medicinal Chemistry (1994), 37(5), 701-5
CODEN: JMCMAR; ISSN: 0022-2623

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A novel series of octapeptide LHRH antagonists was designed based on LHRH agonist H-Phe-Trp-Ser-Tyr-D-Trp-Leu-Arg-Pro-NHEt. By adopting a systematic SAR study, first the in vitro activity was improved, followed by the in vivo LH suppression, raising them up to the range of the decapeptide antagonists NalGlu and A-75998, resulting in antagonist 4-FC6H4CH2CH2CO-D-Nal-Ser-MeTyr-D-Lys(Nic)-Leu-Lys(CHMe2)-Pro-D-Ala-NH2 (Nal = 3-(1-naphthyl)alanine, Nic = nicotinoyl) (A-76154). The octapeptide antagonist A-76154 is the most potent reduced-size LHRH antagonist reported. It suppresses LH in the castrated rat by over 80% for a period of 4 h following s.c. bolus administration of 30 µg/kg.

IT 136988-34-6P


RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and LH releasing hormone antagonist activity of)



RN 136988-34-6 HCAPLUS

CN L-Prolinamide, 4-chloro-N-[3-(4-fluorophenyl)-1-oxopropyl]-D-phenylalanyl-L-seryl-L-tyrosyl-D-tryptophyl-L-leucyl-L-arginyl-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





L11 ANSWER 61 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1993:662666 HCAPLUS

DOCUMENT NUMBER: 119:262666

TITLE: Characterization of cloned somatostatin receptors
SSTR4 and SSTR5

AUTHOR(S): Raynor, Karen; O'Carroll, Anne Marie; Kong, Haeyoung;
Yasuda, Kazuki; Mahana, Lawrence C.; Bell, Graeme I.;
Reisine, Terry

CORPORATE SOURCE: Sch. Med., Univ. Pennsylvania, Philadelphia, PA,
19104, USA

SOURCE: Molecular Pharmacology (1993), 44(2), 385-92
CODEN: MOPMA3; ISSN: 0026-895X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The recent mol. cloning of the genes and cDNAs encoding multiple somatostatin (SRIF) receptor subtypes has allowed for the individual expression of these receptors in mammalian cells and characterization of their resp. pharmacol. profiles. Previously, the authors fully described and compared the pharmacol. properties of the first 3 SRIF receptor subtypes, SRIF receptor type (SSTR)1, SSTR2, and SSTR3. In the present study, the authors have investigated the properties of the newly cloned SRIF receptor subtypes SSTR4 and SSTR5 with regard to pharmacol. profiles, the regulation of high-affinity agonist binding to these receptors by stable GTP analogs, Na⁺, or prior exposure to agonists, and the inhibition of forskolin-stimulated cAMP accumulation mediated by these receptors.

The authors labeled SSTR4 and SSTR5 expressed in Chinese hamster ovary (CHO-K1) and COS-1 cells, resp., with the metabolically stable SRIF analog 125I-CGP 23996. Radioligand binding competition studies were performed using SRIF analogs of differing structures, including hexapeptide analogs similar to MK 678, octapeptide analogs similar to SMS 201-995, pentapeptide analogs similar to c[Ahep-Phe-D-Trp-Lys-Thr(Bzl)], and linear SRIF analogs. SSTR4 bound compds. in all structural classes with high to moderate affinities, and several compds. were identified that are >100-fold selective for SSTR4, compared with the other cloned SRIF receptors, including the linear SRIF analog BIM 23052 and the CGP 23996-like SRIF analog L 362,855. In contrast, SSTR5 bound very few SRIF analogs with high affinity. Both receptors could be regulated by prior exposure to agonist. In addition, agonist binding to SSTR4 was reduced by stable GTP analogs, Na⁺, and pertussis toxin, but agonist binding to SSTR5 was not affected by these treatments. SSTR4 is efficiently coupled to the inhibition of adenylyl cyclase activity, whereas SSTR5 appears not to couple to this cellular effector system. Such differences between the cloned SRIF receptors provide useful strategies for identifying regions of these receptor subtypes that may be involved in ligand-binding specificities and G protein and cellular effector system coupling. The identification of subtype-selective SRIF analogs may lead to more specific therapeutic interventions.

IT 150155-65-0

RL: BIOL (Biological study)


(cloned somatostatin SSTR4 and SSTR5 receptors interaction with)


RN 150155-65-0 HCAPLUS

CN D-Phenylalaninamide, 4-chloro-D-phenylalanyl-L-alanyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





L11 ANSWER 62 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1993:650508 HCAPLUS

DOCUMENT NUMBER: 119:250508

TITLE: Preparation of 5-amino-4-hydroxyhexanoic acid derivative containing peptides as HIV protease inhibitors

INVENTOR(S): Lang, Marc; Bold, Guido; Faessler, Alexander; Schneider, Peter; Van Hoogesvest, Peter

PATENT ASSIGNEE(S): Ciba-Geigy A.-G., Switz.

SOURCE: Eur. Pat. Appl., 79 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 532466	A2	19930317	EP 1992-810678	19920903
EP 532466	A3	19930616		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 05230095	A	19930907	JP 1992-238424	19920907
CA 2077948	A1	19930313	CA 1992-2077948	19920910
AU 9222889	A	19930318	AU 1992-22889	19920910
AU 661018	B2	19950713		
IL 103126	A	19970930	IL 1992-103126	19920910
NO 9203533	A	19930315	NO 1992-3533	19920911
HU 63632	A2	19930928	HU 1992-2925	19920911
ZA 9206938	A	19940311	ZA 1992-6938	19920911
PL 169969	B1	19960930	PL 1992-295905	19920911
RU 2067585	C1	19961010	RU 1992-5052915	19920911
CN 1089269	A	19940713	CN 1993-100044	19930104
PRIORITY APPLN. INFO.:				
			CH 1991-2689	A 19910912
			CH 1992-980	A 19920327



09890219

CH 1992-2007

A 19920625

OTHER SOURCE(S):
GI

MARPAT 119:250508



AB Title compds. [I; R1 = H, alkoxycarbonyl, heterocyclylcarbonyl, heterocyclyloxycarbonyl, (substituted) benzyloxycarbonyl, etc.; X = bond, α -amino acid residue; R2, R3 = (substituted) Ph, cyclohexyl; A1 = bond, α -amino acid residue; A2 = α -amino acid residue; A1A2 = dipeptide residue whose central amide bond is reduced; NR4R5 = (thio)morpholino], were prepared as HIV protease inhibitors. Thus, title compound II was prepared in many steps starting from BOC-phenylalaninal using solution phase methods. I inhibited HIV-1 multiplication in MT-2 cells with ED90's of 10-5-10-8M. Generic I oral formulations are given.

IT 150608-23-4P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation of, as HIV protease inhibitor)

RN 150608-23-4 HCAPLUS

CN Carbamic acid, [1,4-bis[(4-fluorophenyl)methyl]-2-hydroxy-5-[[1-[[[1-[(4-methoxyphenyl)methyl]-2-(4-morpholinyl)-2-oxoethyl]amino]carbonyl]-2-methylpropyl]amino]-5-oxopentyl]-, 1,1-dimethylethyl ester, [1S-[1R*,2R*,4S*,5[R*(R*)]]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search



L11 ANSWER 63 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1993:574313 HCAPLUS

DOCUMENT NUMBER: 119:174313

TITLE: Cloned somatostatin receptors: Identification of subtype-selective peptides and demonstration of high affinity binding of linear peptides

AUTHOR(S): Raynor, Karen; Murphy, William A.; Coy, David H.; Taylor, John E.; Moreau, Jacques Pierre; Yasuda, Kazuki; Bell, Graeme I.; Reisine, Terry

CORPORATE SOURCE: Sch. Med., Univ. Pennsylvania, Philadelphia, PA, 19104, USA

SOURCE: Molecular Pharmacology (1993), 43(6), 838-44
CODEN: MOPMA3; ISSN: 0026-895X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The authors investigated the affinities of a battery of somatostatin (SRIF) analogs to bind to SRIF receptor subtypes SSTR1 (cloned somatostatin complex), SSTR2, and SSTR3, as well as their abilities to inhibit the release of growth hormone from anterior pituitary cells in vitro. SSTR1 and SSTR3 receptors expressed in Chinese hamster ovary and COS-1 cells, resp., were labeled with the metabolically stable SRIF analog 125I-CGP 23996. SSTR2 receptors expressed in Chinese hamster ovary cells were labeled with the SSTR2-specific radioligand 125I-MK-678. Inhibition studies were performed using SRIF analogs of differing structures, including hexapeptide analogs similar to MK-678, octapeptide analogs similar to SMS 201-995, pentapeptide analogs similar to c[Ahep-Phe-D-Trp-Lys-Thr(Bzl)] (SA), and linear SRIF analogs. SSTR1 bound SRIF and SRIF-28 with high affinity and the peptide SA and its structural analogs with low affinity. The hexapeptides did not interact with SSTR1 at concns. as high as 1 μ M, and only a few of the octapeptides or linear peptides bound, with very low affinities. In contrast, 125I-MK-678 binding to SSTR2 was potently inhibited by the hexapeptides, octapeptides, and some of the linear compds., whereas SA and its analogs did not bind to SSTR2. The potencies of the various SRIF agonists to inhibit growth hormone release in vitro was highly correlated with their potencies to inhibit radioligand binding to SSTR2, but not to SSTR1 or SSTR3. SSTR3

09890219

bound analogs of each class but with moderate to low affinities, with the exception of several linear peptides and one of the octapeptides. For the first time the binding affinities of linear analogs of SRIF, some of which display subnanomolar affinities and are highly selective for SRIF receptor subtypes, are reported. Most importantly, these studies identify several peptide analogs that are highly potent, specific, and selective for individual subtypes of SRIF receptors. Such information, coupled with the knowledge of the distribution of these receptor subtypes in normal and pathol. tissues, will be critical for more specific exptl. and therapeutic interventions.

IT 150155-65-0, BIM 23067

RL: BIOL (Biological study)


(somatostatin receptor subtype binding of, selectivity in relation to)

RN 150155-65-0 HCAPLUS


CN D-Phenylalaninamide, 4-chloro-D-phenylalanyl-L-alanyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



Updated Search



L11 ANSWER 64 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1993:423988 HCAPLUS

DOCUMENT NUMBER: 119:23988

TITLE: Stoichiometric labeling of peptides by iodination on tyrosyl or histidyl residues

AUTHOR(S): Tsomides, Theodore J.; Eisen, Herman N.

CORPORATE SOURCE: Cent. Cancer Res., Massachusetts Inst. Technol., Cambridge, MA, 02139, USA

SOURCE: Analytical Biochemistry (1993), 210(1), 129-35
CODEN: ANBCA2; ISSN: 0003-2697

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Radioiodination with ^{125}I or ^{131}I is a favored technique for labeling biol. active peptides or proteins because of high specific radioactivities and convenience in counting γ -emissions. Previous studies used trace labeling, in which fewer than 1% of the mols. are iodinated. Procedures are described for obtaining stoichiometrically iodinated and therefore chemical homogeneous peptides with specific activities exceeding 107 cpm/ μg (≈ 10 Ci/mmol). By analyzing the pH dependence of iodination on tyrosyl and histidyl residues, it is shown that the method described can be applied to many short peptides and optimized for labeling on tyrosine and/or histidine. The power of reverse-phase HPLC is exploited to resolve multiple products substituted with different molar equivalents of iodine from each other and from unlabeled peptide. Specific radioactivity ratios can be used to identify the products, as confirmed by Edman sequence anal. under conditions that separated iodinated tyrosine and histidine derivs. from all other amino acids. It is also shown that the biol. activities of iodinated and uniodinated peptides can differ by several orders of magnitude in a T cell assay and the usefulness of stoichiometric labeling to overcome ambiguities inherent in studying biol. activities with trace-labeled peptides is also demonstrated.

IT 148362-79-2P

RL: PREP (Preparation)
(preparation and characterization of)


RN 148362-79-2 HCAPLUS

09890219


CN L-Leucine, N-[N-[N2-[N-[N-[N-(N-L-arginylglycyl)-3-iodo-L-tyrosyl]-L-valyl]-3-iodo-L-tyrosyl]-L-glutaminy]-L- α -glutamyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



PAGE 1-B



PAGE 2-A



L11 ANSWER 65 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1992:585141 HCAPLUS
DOCUMENT NUMBER: 117:185141
TITLE: Stabilization of the N-terminal residues of
luteinizing hormone-releasing hormone agonists and the

Updated Search

effect on pharmacokinetics

AUTHOR(S): Haviv, Fortuna; Fitzpatrick, Timothy D.; Nichols, Charles J.; Swenson, Rolf E.; Bush, Eugene N.; Diaz, Gilbert; Nguyen, A.; Nellans, Hugh N.; Hoffman, Daniel J.; et al.

CORPORATE SOURCE: Pharm. Prod. Div., Abbott Lab., Abbott Park, IL, 60064, USA

SOURCE: Journal of Medicinal Chemistry (1992), 35(21), 3890-4
CODEN: JMCMAR; ISSN: 0022-2623

DOCUMENT TYPE: Journal

LANGUAGE: English

AB To stabilize leuprolide, [D-Leu6,Pro9-NH₂]LHRH (LHRH = LH-releasing hormone), against chymotrypsin and intestinal degradation, several agonists of LHRH, e.g. [N-Me-Ser4,D-Leu6,Pro9-NH₂]LHRH, [N-Ac-Sar1,D-Leu6,Pro9-NH₂]LHRH (Sar = sarcosine), [Phe2,D-Trp,Pro9-NH₂]LHRH, [N-MePhe2,D-Leu6,Pro9-NH₂]LHRH, [Tyr(Me)3,D-Leu6,Pro9-NH₂]LHRH, modified at positions 1, 2, or 3 and/or containing N- α -Me at positions 1, 2, or 4, were synthesized by the solid-phase method. These agonists were tested in vitro for (a) rat pituitary LHRH receptor binding, (b) LH release from rat pituitary cells, (c) stability against chymotrypsin, and (d) stability against rat intestinal degradation. The clearances of the compounds in the rat were determined using a RIA. Complete stabilization against chymotrypsin (t_{1/2}) and luminal degradation (T_{1/2}) was achieved with substitution of NMe-Ser4 in leuprolide; however, with an increase in clearance. Substitution with 1-Nal3 (Nal = naphthylalanine) increased both t_{1/2} and T_{1/2}, while substitution with NAc-Sar1 increased only T_{1/2}. [NAcSar1,NMeSer4,D-Trp6,Pro9NH₂]LHRH, the doubly stabilized analog, was tested in the rat by both i.v. and id administrations, and its bioavailabilities were measured. No significant improvement in id absorption over leuprolide was observed.


IT 143399-05-7P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation, LH-releasing activity and stability of, to chymotrypsin and intestinal degradation)


RN 143399-05-7 HCAPLUS

CN Luteinizing hormone-releasing factor (swine), 3-(4-chloro-L-phenylalanine)-6-D-leucine-9-(N-ethyl-L-prolinamide)-10-deglycinamide- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A





L11 ANSWER 66 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1992:214921 HCAPLUS
 DOCUMENT NUMBER: 116:214921
 TITLE: Preparation of iodinated peptides
 INVENTOR(S): Halatsch, Wolf Rainer; Sohr, Reinhard; Henklein, Peter; Schmidt, Eberhard
 PATENT ASSIGNEE(S): Humboldt-Universitaet zu Berlin, Germany
 SOURCE: Ger. (East), 4 pp.
 CODEN: GEXXA8
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DD 296083	A5	19911121	DD 1990-339759	19900412
DD 296083	B5	19970220		
PRIORITY APPLN. INFO.:			DD 1990-339759	19900412
OTHER SOURCE(S):	CASREACT 116:214921; MARPAT 116:214921			
GI				



AB Title compds. were prepared via reacting free amino group-containing peptides H₂N-AS₁...AS_n-B (I; AS₁-AS_n = any amino acid residue; A, B = protecting group or any organic group) or A-NH-AS₁...A(NH₂)S_x-B (II; AS_x = amino acid residue containing an NH₂ group in the sidechain) with norborn-5-ene-2,3-carboxyimide III (R = I, I125) or acylating I or II with III (R = H) and subsequent iodination. III (R = H) (preparation given) in dioxane containing Zn

09890219

was treated with KI in an NaOAc buffer, Chloramine T in NH₄OAc was added to the reaction mixture, and the resulting mixture was stirred for 30 s to give III (R = I). This was condensed with H-Asp-Tyr(SO₃Na)-Met-Gly-Trp-Met-Asp-Phe-NH₂ in DMF-pyridine containing (Me₂CH)₂NEt to give Q-Asp-Tyr(SO₃Na)-Met-Gly-Trp-Met-Asp-Phe-NH₂ [Q = 4,3-(HO)IC₆H₃CH₂CH₂CO].

IT 140908-81-2P

RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation of, as pharmaceutical and diagnostic agent)

RN 140908-81-2 HCAPLUS


CN Caerulein, 1-de(5-oxo-L-proline)-2-de-L-glutamine-3-[N-[3-(4-hydroxy-3-iodophenyl)-1-oxopropyl]-L-aspartic acid]-5-L-methionine-, monosodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



● Na



L11 ANSWER 67 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1992:129629 HCAPLUS
 DOCUMENT NUMBER: 116:129629
 TITLE: Preparation of reduced size LH-RH analogs as LH-RH agonists and antagonists
 INVENTOR(S): Haviv, Fortuna; Palabrica, Christopher A.; Greer, Jonathan; Fitzpatrick, Timothy D.
 PATENT ASSIGNEE(S): Abbott Laboratories, USA
 SOURCE: Eur. Pat. Appl., 90 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 417454	A2	19910320	EP 1990-114752	19900801
EP 417454	A3	19910710		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
US 5140009	A	19920818	US 1990-548511	19900710
CA 2022437	A1	19910208	CA 1990-2022437	19900801
CA 2022437	C	20021022		
NO 9003454	A	19910208	NO 1990-3454	19900806
HU 55414	A2	19910528	HU 1990-4911	19900806
KR 161972	B1	19981116	KR 1990-11998	19900806
AU 9060285	A	19910207	AU 1990-60285	19900807
JP 03081292	A	19910405	JP 1990-209059	19900807
AU 9457894	A	19940519	AU 1994-57894	19940317
AU 675274	B2	19970130		
PRIORITY APPLN. INFO.:			US 1989-390269	A 19890807
			US 1990-548511	A 19900710
			US 1988-154682	B2 19880210
OTHER SOURCE(S):		MARPAT 116:129629		

AB Reduced size LH-RH analogs T-Q-X-A-B-C-D-E-F-Y [T = absent, D- or L-H-Gln(Et), Z-W-W1CO; Z = H, C1-6 alkyl, cycloalkyl, etc.; W = absent, alkylene, alkenylene; W1 = absent, O, S, NH; Q = absent, (substituted) D- or L-Phe, His, Trp, etc.; X = absent, (substituted) D- or L-Trp, 3-(1-naphthyl)alanyl, Pro, etc.; A = (substituted) L-Ser, Ala, Gln, etc.; B = (substituted) Tyr, Trp, His, etc.; C = (substituted) D-amino acid residue, Ser(PO3H2), Ser(PO3Me2), etc.; D = (substituted) Leu, Ile, Thr(PO3H2), etc.; E = L-amino acyl residue NR1CH[(CH2)pR2]CO, etc.; R1 = H, Me, Et, Pr, Me2CH; R2 = NH2, alkylamino, cycloalkylamino, alkanoylamino, etc.; p = 1-4; F = L-Pro, trans- β -aminocyclopentanecarbonyl, etc.; Y = D- or L-Ala-NH2, Gly-NH2, etc.; with provisos] were prepared. Thus, 1-naphthylacetyl-Ser-Tyr-D-Leu-Leu-Arg-Pro-NHEt (I) was prepared via solid phase methods starting with resin-bound Boc-Pro-OH and Boc-Arg(Tos)-OH, Boc-Leu-OH, Boc-D-Leu-OH, Boc-Tyr(4-BrZ)-OH, Boc-Ser(Bzl)-OH, and naphthylacetic acid. I had a pD2 (neg. log of concentration which produces half-maximal release of LH) of 6.85

vs. 9.27 for LH-RH.


IT 136967-92-5P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of, as LH-RH agonist and antagonist)


RN 136967-92-5 HCAPLUS

CN L-Prolinamide, N-[3-(4-fluorophenyl)-1-oxopropyl]-L-seryl-L-tyrosyl-D-tryptophyl-L-leucyl-L-arginyl-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





L11 ANSWER 68 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1991:157334 HCAPLUS

DOCUMENT NUMBER: 114:157334

DOCUMENT NUMBER: 114104551
TITLE: Increased concentrations of immunoreactive inhibin during conception cycles in the marmoset monkey: suppression with an LHRH antagonist and cloprostenol

AUTHOR(S): Webley, G. E.; Knight, P. G.; Given, A.; Hodges, J. K.
CORPORATE SOURCE: Comp. Physiol. Group, Inst. Zool., London, NW1 4RY, UK
SOURCE: Journal of Endocrinology (1991), 128(3), 465-73

CODEN: JOENAK; ISSN: 0022-0795

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Peripheral concns. of immunoreactive (ir) inhibin have been measured during the ovarian cycle and early pregnancy in the marmoset monkey. Blood samples were taken (3/wk) during conception and non-conception cycles. Ir-inhibin was measured by RIA using an antiserum raised against a synthetic peptide fragment of the α subunit of human inhibin. Monomeric bovine α subunit and 32 kDa bovine inhibin were used as tracer and standard resp. In all animals low concns. of ir-inhibin were recorded during the follicular phase (40-60 $\mu\text{g/L}$) of the cycle. After ovulation, ir-inhibin concns. increased but the peak concns. attained differed between conception and non-conception cycles. In non-pregnant animals ir-inhibin concns. reached a maximum of 242 $\mu\text{g/L}$ on days 12/13 after ovulation. In pregnant animals ir-inhibin concns. were higher (1.8-fold) than in non-pregnant animals on days 8/9 after ovulation, and reached a maximum value of 636 $\mu\text{g/L}$ on days 20/21 after ovulation. Administration of an LH-RH antagonist during the luteal phase on days 6-8 after ovulation decreased progesterone and ir-inhibin concns. within 4 and 8 h, resp. This was prevented by coadministration with human chorionic gonadotropin. Administration of cloprostenol to pregnant animals between days 17 and 20 after ovulation halved the initial concns. of both inhibin and progesterone within 1.5 h. The increase in plasma ir-inhibin concns. in the luteal phase and the apparent similarity in control of ir-inhibin and progesterone supports a luteal source of ir-inhibin in both conception and non-conception cycles. The higher levels of ir-inhibin from days 8/9 after ovulation in conception cycles were not related to any detectable increase in peripheral concns. of chorionic gonadotropin and occurred at least 4 days before the expected time of implantation. This suggests a role for the conceptus in inhibin secretion which may involve the release of an embryo message before implantation.

IT 132998-39-1

RL: BIOL (Biological study)

(inhibin secretion suppression by, in marmoset monkey)


09890219

RN 132998-39-1 HCAPLUS


CN D-Alaninamide, N-acetyl-3-(2-naphthalenyl)-D-alanyl-L-histidyl-4-chloro-D-phenylalanyl-L-seryl-L-tyrosyl-D-arginyl-L-phenylalanyl-L-arginyl-L-prolyl-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L11 ANSWER 69 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1990:199087 HCAPLUS

DOCUMENT NUMBER: 112:199087

TITLE: Synthetic studies on physiologically active oligopeptides carrying isodityrosine units

AUTHOR(S): Suzuki, Y.; Nishiyama, S.; Yamamura, S.

CORPORATE SOURCE: Fac. Sci. Technol., Keio Univ., Yokohama, Japan

SOURCE: Tennen Yuki Kagobutsu Toronkai Koen Yoshishu (1989), 31st, 190-6

CODEN: TYKYDS

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

Updated Search

09890219

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB In connection with synthetic studies on physiol. active natural products carrying isodityrosine units, the total synthesis of OF4949-III (I), and K-13 (II) along with a synthetic study on vancomycin have been achieved. I and II have been successfully synthesized from the corresponding tripeptides. Interesting activities against gram pos. bacteria and the concepts of mol. recognition have evoked total synthesis of vancomycin. Methods used to construct the isotyrosine units in this antibiotic are discussed. I is an aminopeptidase B inhibitor and II is an inhibitor of angiotensin-converting enzyme.


IT 123418-38-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and oxidation-cyclization of)

RN 123418-38-2 HCAPLUS

CN L-Tyrosine, N-[N-(N-acetyl-3,5-dichloro-L-tyrosyl)-O-(phenylmethyl)-L-tyrosyl]-3,5-diiodo-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 70 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1990:77969 HCAPLUS

DOCUMENT NUMBER: 112:77969

TITLE: Preparation of antibiotic OF4949 analogs

INVENTOR(S): Yamamura, Shosuke; Nishiyama, Shigeru; Suzuki, Ryoichi; Katayama, Kaoru

PATENT ASSIGNEE(S): Takara Shuzo Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 10 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 01168698	A	19890704	JP 1987-325494	19871224

Updated Search

09890219

PRIORITY APPLN. INFO.:


JP 1987-325494

19871224

OTHER SOURCE(S):

MARPAT 112:77969

GI



AB The title compds. [I; R1 = H, alkyl, acyl; R2 = H, OH, acyloxy; R3 = H, alkyl, protecting group; R4 = H, alkyl, protecting group; X = H2NCO, hydroxyalkyl, CO2H, alkoxycarbonyl; Y = H2NCO, hydroxyalkyl, (protected) CO2H; Z = H] (no data on pharmacol. activities), are prepared via catalytic reduction of I (Z = Br; R1-R4, X, and Y same as defined above). I (R1 = Me, X = H2NCO, R2 = R3 = H, R4 = Me3CO2C, Y = CO2Me, Z = Br) in MeOH was hydrogenolyzed over Pd black to give 96% I (R1 = Me, X = H2NCO, R2 = R3 = H, R4 = Me3CO2C, Y = CO2Me, Z = H).


IT 116523-66-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, as intermediate for antibiotics)

RN 116523-66-1 HCAPLUS

CN L-Tyrosine, 3,5-dichloro-N-[N2-[3,5-dibromo-N-[(phenylmethoxy)carbonyl]-L-tyrosyl]-L-asparaginyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 71 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1989:595381 HCAPLUS

DOCUMENT NUMBER: 111:195381

Updated Search

09890219

TITLE: Biomimetic synthesis and stereostructure of K-13, a novel inhibitor of angiotensin I converting enzyme
AUTHOR(S): Nishiyama, Shigeru; Suzuki, Yoshikazu; Yamamura, Shosuke
CORPORATE SOURCE: Fac. Sci. Technol., Keio Univ., Yokohama, Japan
SOURCE: Tetrahedron Letters (1989), 30(3), 379-82
CODEN: TELEAY; ISSN: 0040-4039

DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 111:195381


AB A novel inhibitor of angiotensin I converting enzyme, K-13, has been synthesized from N-acetyl-3,5-dichloro-L-tyrosyl-O-benzyl-L-tyrosyl-3,5-diiodo-L-tyrosine Me ester, whose oxidation with thallium trinitrate as a key step followed by zinc reduction affords the corresponding di-Ph ether with the same heterocyclic skeleton as that of K-13, indicating that K-13 is biosynthesized from three mols. of L-tyrosine.

IT 123418-33-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and oxidative cyclization of, with thallium trinitrate)

RN 123418-33-7 HCAPLUS

CN L-Tyrosine, 3,5-dibromo-N-[N-[3,5-dichloro-N-[(1,1-dimethylethoxy)carbonyl]-L-tyrosyl]-O-(phenylmethyl)-L-tyrosyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 72 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1989:433726 HCAPLUS

DOCUMENT NUMBER: 111:33726

TITLE: Synthesis of tritium labeled derivatives of atrial natriuretic peptide (ANP) and characterization of a biologically active linear analog

AUTHOR(S): Pham, P.; Moustier, A. M.; Rousseau, B.; Beaucourt, J. P.

CORPORATE SOURCE: Serv. Mol. Marquees, CEN-Saclay, Gif-sur-Yvette, 91191, Fr.

SOURCE: Colloque INSERM (1989), 174(Forum Pept., 2nd, 1988), 247-50

CODEN: CINMDE; ISSN: 0768-3154

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The syntheses of three tritiated rat ANP(1-28) derivs.: (4-3H-Phe8)ANP,

Updated Search

09890219

(4-3H-Phe26)ANP and (4-3H-Phe8, 3,5-3H2-Tyr28)ANP are described. The high specific activity peptides obtained are biol. fully active. A byproduct of the tritiation reaction, isolated by HPLC, was characterized as the linear analog of ANP: (Ala7,23)ANP. Preliminary results showed this peptide to be biol. active and suggest that the disulfide bridge of ANP is not essential for activity.

IT 120642-96-8

RL: RCT (Reactant); RACT (Reactant or reagent)
(tritiation of)

RN 120642-96-8 HCAPLUS

CN Atrial natriuretic peptide-28 (rat), 26-(4-iodo-L-phenylalanine)- (9CI)
(CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

L11 ANSWER 73 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1989:232086 HCAPLUS

DOCUMENT NUMBER: 110:232086

TITLE: Preparation of fluorine-containing atrial natriuretic peptides as diuretics, natiuretics, and antihypertensives

INVENTOR(S): Rakhit, Sumanas; Goghari, Mahesh H.

PATENT ASSIGNEE(S): Bio-Mega Inc., Can.

SOURCE: Eur. Pat. Appl., 16 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
EP 283956	A2	19880928	EP 1988-104365	19880318
EP 283956	A3	19900411		
EP 283956	B1	19921111		
R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
IL 85728	A	19930114	IL 1988-85728	19880314
AU 8813167	A	19881124	AU 1988-13167	19880316
AU 605259	B2	19910110		
AT 82295	T	19921115	AT 1988-104365	19880318
JP 01006296	A	19890110	JP 1988-71683	19880325
PRIORITY APPLN. INFO.:				CA 1987-532982 A 19870325
				CA 1987-542192 A 19870715
				EP 1988-104365 A 19880318

OTHER SOURCE(S): MARPAT 110:232086

GI

Y-R1-R2-Gly-Arg-R3-Asp-Arg-Ile-Gly-

Ala-Gln-Ser-Gly-Leu-Gly-Cys-Asn-Ser-R4-

Arg-R5-Z

I

AB The title compds. (I; R1, R4 = Phe, 2-, 4-, or 5-fluorophenylalanyl, trisfluorophenylalanyl; R2 = Gly, Ala, D-Ala; R3 = Ile, Met; R5 = Tyr,

Updated Search

09890219

Null; Y = thioalkylcarbonyl, R6 - Cys; R6 = H-Ser-Ser, H-Arg-Ser, H-Arg-Arg-Ser-Ser, H-Leu-Arg-Arg-Ser-Ser, H-Ser-Leu-Arg-Arg-Ser-Ser; Z = OH, amino) useful as diuretics, natriuretics, and antihypertensives; were prepared I (Y = H-Ser-Leu-Arg-Ser-Ser-Cys, R1 = Phe, R2 = Gly, R3 = Met, R4 = 4-fluorophenylalanyl, R5 = Tyr, Z = OH), prepared on (α -phenylacetamido)benzylbenzhydrylamine resin, had a relative potency of 0.94 v.s. human atrial natriuretic peptide (hANP) in the rabbit aorta assay.

IT 120728-19-0P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation of, as atrial natriuretic peptide analog)
RN 120728-19-0 HCAPLUS
CN Atrial natriuretic peptide-28 (human), 26-(4-fluoro-L-phenylalanine)-(9CI) (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

L11 ANSWER 74 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1989:213362 HCAPLUS
DOCUMENT NUMBER: 110:213362
TITLE: Preparation of cyclic peptides as anticancer agents
INVENTOR(S): Itokawa, Hideji; Watanabe, Kinzo; Kawaoto, Satoshi; Inoue, Tsutomu
PATENT ASSIGNEE(S): Tobishi Pharmaceutical Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 10 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
JP 63203671	A	19880823	JP 1987-34508	19870219
PRIORITY APPLN. INFO.:			JP 1987-34508	19870219
OTHER SOURCE(S):	MARPAT	110:213362		

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I (R1 = H, alkyl; R2 = H, PhCH2O2C; R3 = H, (substituted) alkyl; X-X2 = halo) useful as intermediates for anticancer agents I (X-X2 = H), are prepared from tripeptides II (R4 = Me; X3 = halo) or their amino- or CO2R4-protected compds. via cyclic tripeptides III. Treatment of II (X-X3 = Br; R2 = PhCH2O2C; R3 = CH2CHMe2; R4 = Me) (preparation given) with thallium nitrate in MeOH and reduction of a product, after work-up and chromatog., with Zn in AcOH gave the Me ester of I (X-X2 = Br; R1 = H; R2 = PhCH2O2C; R3 = CH2CHMe2) (IV), which showed 6.0 μ g/mL of IC50 against P388 mice leukemia cells. IV in MeOH was hydrogenated in the presence of KOAc and Pd/C to give I (X-X2 = H; R1 = H; R2 = PhCH2O2C; R3 = CH2CHMe2).

IT 120377-40-4P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and reaction of, in preparation of cyclic peptide anticancer agent)


Updated Search

09890219

RN 120377-40-4 HCAPLUS

CN L-Tyrosine, 3,5-dibromo-N-[N-[3,5-dibromo-N-[(phenylmethoxy)carbonyl]-L-tyrosyl]-L-leucyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 75 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1989:213307 HCAPLUS

DOCUMENT NUMBER: 110:213307

TITLE: Synthesis of tritium labelled atrial natriuretic factor (ANF) derivatives and characterization of a tritiated biologically active linear peptide by-product

AUTHOR(S): Pham, P.; Moustier, A.; Rousseau, B.; Beaucourt, J. P.
CORPORATE SOURCE: Serv. Mol. Marquees, CEN-Saclay, Gif-sur-Yvette, 91191, Fr.

SOURCE: Journal of Labelled Compounds and Radiopharmaceuticals (1988), 25(8), 901-11
CODEN: JLCRD4; ISSN: 0362-4803

DOCUMENT TYPE: Journal

LANGUAGE: French

OTHER SOURCE(S): CASREACT 110:213307

AB (4-3H-Phe8)ANF, (4-3H-Phe26)ANF, and (4-3H-Phe8, 3,5-3H2-Tyr28)ANF were obtained by catalytic dehalogenation of iodinated precursors with 3H-Pd and exhibited high specific radioactivity. A radioactive byproduct isolated by HPLC was characterized as 3H-(Ala7,23)ANF and was biol. active.

IT 120642-96-8

RL: RCT (Reactant); RACT (Reactant or reagent)
(reductive tritiation of)

RN 120642-96-8 HCAPLUS

CN Atrial natriuretic peptide-28 (rat), 26-(4-iodo-L-phenylalanine)- (9CI)
(CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

L11 ANSWER 76 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:611493 HCAPLUS

DOCUMENT NUMBER: 109:211493

TITLE: Preparation of somatostatin analogs as drugs

INVENTOR(S): Coy, David H.; Murphy, William A.; Heiman, Mark L.

PATENT ASSIGNEE(S): Tulane Educational Fund, Inc., USA

SOURCE: Eur. Pat. Appl., 5 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

Updated Search

09890219

LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 5
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 277419	A2	19880810	EP 1987-310487	19871127
EP 277419	A3	19900214		
EP 277419	B1	19970618		
R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 63196599	A	19880815	JP 1987-295911	19871124
JP 2568228	B2	19961225		
AT 154612	T	19970715	AT 1987-310487	19871127
ES 2104551	T3	19971016	ES 1987-310487	19871127
US 4853371	A	19890801	US 1988-209883	19880622
US 4904642	A	19900227	US 1989-312138	19890217
PRIORITY APPLN. INFO.:			US 1987-10349	A 19870203
			US 1985-775488	A2 19850912
			US 1986-875266	A2 19860617
			US 1987-70400	A2 19870707
			US 1988-209883	A3 19880622

OTHER SOURCE(S): MARPAT 109:211493

AB R-A1-Cys-Tyr-D-Trp-Lys-A2-Cys-A3 (I; R = H, C1-20 alkyl; A1 = D- β -Nal, D-Trp, D-X-Phe; A2 = α -aminobutyryl; A3 = Thr-NH₂, Thr-OH, Nal-NH₂, Trp-NH₂; X = H, OH, Me, halo) and pharmaceutically acceptable salts thereof were prepared for reducing growth hormone, insulin, glucagon, and/or pancreatic exocrine secretion. D- β -Naphthylalanyl-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂ was prepared by the solid-phase method using BOC-protected amino acids on benzhydrylamine resin.

IT 117382-74-8P


RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as somatostatin analog)

RN 117382-74-8 HCAPLUS


CN L-Threoninamide, 4-chloro-D-phenylalanyl-L-cysteinyl-L-tyrosyl-D-tryptophyl-L-lysyl-2-aminobutanoyl-L-cysteinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A



Updated Search



L11 ANSWER 77 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1988:529660 HCAPLUS
DOCUMENT NUMBER: 109:129660
TITLE: Total synthesis of OF4949-III, a novel inhibitor of
aminopeptidase B
AUTHOR(S): Nishiyama, Shigeru; Suzuki, Yoshikazu; Yamamura,
Shosuke
CORPORATE SOURCE: Fac. Sci. Technol., Keio Univ., Hiyoshi, Japan
SOURCE: Tetrahedron Letters (1988), 29(5), 559-62
CODEN: TELEAY; ISSN: 0040-4039
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 109:129660
GI



AB OF4949-III (I) has been synthesized from N-benzyloxycarbonyl-3,5-dibromo-L-tyrosyl-L-asparaginyl-3,4-dichloro-L-tyrosine Me ester, whose oxidation with

09890219

thallium trinitrate as a key step followed by zinc reduction affords the corresponding di-Ph ether with the same heterocyclic skeleton as that of I.


IT 116523-66-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and oxidative cyclization of)

RN 116523-66-1 HCAPLUS

CN L-Tyrosine, 3,5-dichloro-N-[N2-[3,5-dibromo-N-[(phenylmethoxy)carbonyl]-L-tyrosyl]-L-asparaginyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 78 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:132324 HCAPLUS

DOCUMENT NUMBER: 108:132324

TITLE: Preparation of somatostatin analogs as drugs

INVENTOR(S): Coy, David H.; Murphy, William A.; Heiman, Mark L.

PATENT ASSIGNEE(S): Tulane Educational Fund, Inc., USA

SOURCE: Eur. Pat. Appl., 13 pp.

CODEN: EPXXDW

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 214872	A2	19870318	EP 1986-307044	19860912
EP 214872	A3	19890906		
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
AU 8662076	A	19870319	AU 1986-62076	19860829
AU 602657	B2	19901025		
DK 8604351	A	19870313	DK 1986-4351	19860911
DK 172212	B1	19980105		
FI 8603680	A	19870313	FI 1986-3680	19860911
FI 89062	B	19930430		
FI 89062	C	19930810		
NO 8603638	A	19870313	NO 1986-3638	19860911
NO 174809	B	19940405		
NO 174809	C	19940713		
ES 2003739	A6	19881116	ES 1986-1814	19860911

Updated Search

09890219

CA 1338301	C	19960430	CA 1986-517969	19860911
JP 62116594	A	19870528	JP 1986-215581	19860912
JP 2563278	B2	19961211		
US 4853371	A	19890801	US 1988-209883	19880622
US 4904642	A	19900227	US 1989-312138	19890217
PRIORITY APPLN. INFO.:			US 1985-775488	A 19850912
			US 1986-875266	A 19860617
			US 1987-10349	A2 19870203
			US 1987-70400	A2 19870707
			US 1988-209883	A3 19880622

GI

A1A2NCHA³CO-Cys-A⁴-D-Trp-Lys-A⁵-Cys-A⁷-NH₂ I

AB The title compds. [I; A1, A2 = H, alkyl, phenylalkyl, acyl, alkoxy carbonyl; A3 = CHA6 (A6 = pentafluorophenyl, naphthyl, pyridyl, phenyl); A4 = o-, m-, or p-substituted X-Phe (X = H, halo, NO₂, OH, NH₂, alkyl), pentafluoro-Phe, β -naphthylalanyl (β -Nal); A5 = Thr, Ser, Phe, Val, α -aminoisobutyric acid residue, Ile; A7 = Thr, Trp, β -Nal], somatostatin analogs, and their pharmaceutically acceptable salts are prepared via the solid-phase method. H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂ was prepared by peptide coupling of the appropriate protected amino acids on a benzhydrylamine resin, followed by deprotection and resin cleavage using HF/anisole.



IT 113294-83-0P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, for reduction of growth hormone, insulin, glucagon, or pancreatic exocrine secretion)

RN 113294-83-0 HCAPLUS

CN L-Threoninamide, 2,3,4,5,6-pentafluoro-D-phenylalanyl-L-cysteinyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-cysteinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 79 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1987:214405 HCAPLUS

DOCUMENT NUMBER: 106:214405

TITLE: Somatostatin octapeptide analogs with growth hormone

09890219

release-inhibiting activity
 INVENTOR(S): Coy, David H.; Murphy, William A.; Heiman, Mark L.
 PATENT ASSIGNEE(S): Tulane Educational Fund, Inc., USA
 SOURCE: Eur. Pat. Appl., 12 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 5
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 215171	A2	19870325	EP 1985-308770	19851202
EP 215171	A3	19881109		
EP 215171	B1	19901122		
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
JP 62061997	A	19870318	JP 1985-270577	19851130
JP 08005913	B	19960124		
AT 58542	T	19901215	AT 1985-308770	19851202
US 4853371	A	19890801	US 1988-209883	19880622
US 4904642	A	19900227	US 1989-312138	19890217
PRIORITY APPLN. INFO.:			US 1985-775488	A 19850912
			EP 1985-308770	A 19851202
			US 1986-875266	A2 19860617
			US 1987-10349	A2 19870203
			US 1987-70400	A2 19870707
			US 1988-209883	A3 19880622

GI

RR1NCHR2CO-Cys-Z-D-Trp-Lys-Z1-Cys-Thr-NH2

I

AB The title peptides [I; R, R1 = H, (phenyl)alkyl, COR3, CO2R4; R2 = CH2R5; R3 = alkyl, alkenyl, alkynyl, Ph, naphthyl, phenylalkyl; R4 = (phenyl)alkyl; R5 = pentafluoronaphthyl, pyridyl, Ph, halophenyl, NO2, NH2, OH, alkyl, alkoxy; Z = (un)substituted Ph, pentafluoroalanine, naphthylalanine; Z1 = Thr, Ser, Phe, Val, Ile] or their pharmaceutically acceptable salts, inhibiting the secretion of growth hormone, insulin, and glucagon, were prepared by solid-phase synthesis. I (RR1NCHR2 = N-tert-butoxycarbonyl-D-p-Cl-Phe, Z = Tyr, Z1 = Val) is prepared by using N-tert-butoxycarbonyl-O-benzylthreonine bound to a benzhydrylamine resin.



IT 108335-13-3DP, benzhydrylamine resin-bound
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and deprotection and resin cleavage of)


RN 108335-13-3 HCAPLUS

CN L-Threoninamide, 4-chloro-N-[(1,1-dimethylethoxy)carbonyl]-D-phenylalanyl-S-[(4-methylphenyl)methyl]-L-cysteinyl-L-tyrosyl-D-tryptophyl-N6-[(phenylmethoxy)carbonyl]-L-lysyl-L-valyl-S-[(4-methylphenyl)methyl]-L-cysteinyl-O-(phenylmethyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Updated Search





L11 ANSWER 80 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1987:113675 HCAPLUS

DOCUMENT NUMBER: 106:113675

TITLE: Structure-activity studies of antagonists of luteinizing hormone-releasing hormone with emphasis on the amino-terminal region

AUTHOR(S): Hocart, Simon J.; Nekola, Mary V.; Coy, David H.
CORPORATE SOURCE: Sch. Med., Tulane Univ., New Orleans, LA, 70112, USA
SOURCE: Journal of Medicinal Chemistry (1987), 30(4), 735-9

CODEN: JMCMAR; ISSN: 0022-2623

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The structure-activity relations of the hydrophobic N-terminal region of the antagonist [N-acetyl-D-naphthylalanine, D-p-chlorophenylalanine2,D-Trp3,D-Arg6,Phe7,D-Ala10]-LH-RH [96394-82-0] was investigated by the incorporation of a variety of amino acids with emphasis on positions 1, 2, and 3. The analogs were prepared by routine solid-phase peptide synthesis. All purifications were performed in 2 stages: gel permeation chromatog. followed by preparative, reversed-phase, HPLC. The analogs were assayed in a standard rat antioviulatory assay with a 40% propane-1,2-diol-saline vehicle. A simplified antagonist was developed that allowed the removal of the custom-synthesized D-p-chlorophenylalanine and the labile D-tryptophan while retaining antioviulatory potency. [N-Acetyl-D-naphthylaminel,D-Phe2,3,D-Arg6,Phe7,D-Ala10]-LH-RH [106881-55-4] caused a 56% blockade of ovulation at the 500 ng dose and was approx. equipotent with the parent analog in this system.



IT 106881-64-5

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(ovulation inhibition by, structure in relation to)

RN 106881-64-5 HCAPLUS

CN D-Alaninamide, N-acetyl-3-(2-naphthalenyl)-D-alanyl-3-(2-naphthalenyl)-D-alanyl-2,3,4,5,6-pentafluoro-D-phenylalanyl-L-seryl-L-tyrosyl-D-arginyl-L-phenylalanyl-L-arginyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.





L11 ANSWER 81 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1987:81074 HCAPLUS

DOCUMENT NUMBER: 106:81074

TITLE: Mechanism studies of Coomassie Blue and silver staining of proteins

AUTHOR(S): De Moreno, Miriam R.; Smith, Jean F.; Smith, Robert V.

CORPORATE SOURCE: Coll. Pharm., Univ. Texas, Austin, TX, 78712-1074, USA

SOURCE: Journal of Pharmaceutical Sciences (1986), 75(9), 907-11

CODEN: JPMSAE; ISSN: 0022-3549

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A relatively high complexation affinity has been found for Coomassie Blue G-250 and the following amino acids: arginine; tyrosine; lysine; and histidine. A linear relationship was observed between log molar absorptivity and log mol. weight of 52 and 69 proteins, polypeptides, and di- and tripeptides that were allowed to react with Coomassie Blue G-250 in solution. The solution complexation results were used in a study of the detection of the following model proteins: bovine serum albumin, lysozyme, recombinant DNA derived human insulin, and calmodulin. Interactions between Coomassie Blue stained gels and Ag detection reagents were determined and used as the basis for studies of enhanced sensitivity of detection of electrophoretically developed proteins. Sensitivity enhancements of up to 8-fold were observed when various sulfonic acid dye complexed proteins were detected with Ag reagents vs. the use of Ag reagents alone. A site-directed nucleation of Ag caused by the protein complexed sulfonic acid dyes is proposed as a mechanism for the observed enhancements.

IT 106814-81-7

RL: ANST (Analytical study)


(complexation of, with Coomassie Blue G-250, in solution)

RN 106814-81-7 HCAPLUS


CN D-Alaninamide, N-acetyl-3-(2-naphthalenyl)-D-alanyl-4-chloro-D-phenylalanyl-O-(1-oxohexadecyl)-D-seryl-L-tyrosyl-D-arginyl-L-lysyl-L-arginyl-L-prolyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



Updated Search



L11 ANSWER 82 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1986:627314 HCAPLUS

DOCUMENT NUMBER: 105:227314

TITLE: Analgesic and antipsychotic penta- and heptapeptides

INVENTOR(S): Cervini, Maria Antonietta; De Castiglione, Roberto;

Mena, Renzo; Perseo, Giuseppe; Rossi, Alessandro

PATENT ASSIGNEE(S): Farmitalia Carlo Erba S.p.A., Italy

SOURCE: Ger. Offen.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3537405	A1	19860430	DE 1985-3537405	19851021
GB 2166139	A	19860430	GB 1984-27027	19841025
GB 2166139	B	19880622		
JP 61103898	A	19860522	JP 1985-233569	19851021
BE 903493	A1	19860422	BE 1985-215758	19851022
			GB 1984-27027	A 19841025

PRIORITY APPLN. INFO.:

OTHER SOURCE(S): CASREACT 105:227314; MARPAT 105:227314

GI

Y
|
X-Tyr-A-B-C-E-F-NH₂ I

AB The title peptides [I; X = H, C(:NH)NH₂, amino protective group; Y = H, phenolic OH protective group; A = D-Ala, D-Val, D-Ile, D-Leu, D-Pro, D-Ser, D-Thr, D-Met, D-Met(O), D-Arg, D-Lys, D-Orn; B = Trp, Phg (phenylglycine), p-(un)substituted Phe; C = Phg, Npg (neopentylglycine), Gly, Sar (sarcosine), D- or L-Ala, Val, Ile, Leu, Met, Ser, Thr, Phe, Trp,

Updated Search

09890219

Tyr, etc.; E = D- or L-Tyr, Ser, Thr, Met, Met(O), Leu, Nle, Ape (2-aminovaleric acid), p-(un)substituted Phe; F = Pro-Ser, bond] were prepared as analgesics and antipsychotics. Thus, Boc-Phe-OH was coupled with D-Ala-OBzl.HCl to give Boc-Phe-D-Ala-OBzl, which was converted in 8 steps to H-Tyr-D-Ala-Phe-D-Ala-Tyr-Pro-Ser-NH₂.HCl (II). In the rat tail-flick test II had an ED₅₀ of 1.3 mg/kg s.c.

IT 105412-73-5P


RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, as analgesic and antipsychotic)

RN 105412-73-5 HCAPLUS


CN Dermorphin, 3-(4-chloro-L-phenylalanine)-4-D-alanine- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L11 ANSWER 83 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1986:472825 HCAPLUS

DOCUMENT NUMBER: 105:72825

TITLE: Synthesis and biological activity of highly potent octapeptide analogs of somatostatin

AUTHOR(S): Cai, R. Z.; Szoke, B.; Lu, R.; Fu, D.; Redding, T. W.; Schally, A. V.

Updated Search

09890219



CORPORATE SOURCE: Sch. Med., Tulane Univ., New Orleans, LA, 70146, USA
SOURCE: Proceedings of the National Academy of Sciences of the
United States of America (1986), 83(6), 1896-900
CODEN: PNASA6; ISSN: 0027-8424

DOCUMENT TYPE: Journal
LANGUAGE: English

AB In the search for selective and long-acting analogs of somatostatin, nearly 200 compds. were synthesized by solid-phase methods, purified, and tested biol. Among these octapeptides, some contained N-terminal D-Phe, Ac-D-Phe, or AcPhe followed by hexapeptide sequences Cys-Phe-D-Trp-Lys-Thr-Cys or Cys-Tyr-D-Trp-Lys-Val-Cys and Thr-NH₂ or Trp-NH₂ as C-terminal residues. (Cyclo 2-7)-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂ (I) [99660-13-6] and (cyclo 2-7)-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂ (II) [103222-11-3] were 177 times and 113 times more potent, resp., than somatostatin in tests for inhibition of growth hormone [9002-72-6] release. These 2 octapeptides containing tyrosine and valine in positions 3 and 6, resp., were more active and more selective than their Ph-3 and Thr-6 counterparts, (cyclo 2-7)-D-Phe-Cys-Phe-D-Trp-Lys-thr-Cys-Thr-NH₂ [99685-66-2] and (cyclo 2-7)-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Trp-NH₂ [103222-10-2]. I was also .apprx.6 times more potent than its L-Trp-4 diastereoisomer [103222-07-7]. The analogs I, and II showed a prolonged duration of action and inhibited growth hormone release for at least 3 h. Analogs of both Phe-3/Thr-6 and Tyr-3/Val-6 classes also suppressed the release of insulin [9004-10-8] and glucagon [9007-92-5] in rats and pentagastrin-induced secretion of gastric acid in dogs, but their potencies in these tests were much smaller than the growth-hormone-release inhibitory activity. Some of these analogs possessed antitumor activities as shown by the inhibition of growth of animal models of prostate, mammary, and ductal pancreatic tumors.

IT 103548-89-6
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(growth hormone secretion inhibition by, mol. structure in relation to)
RN 103548-89-6 HCAPLUS
CN L-Threoninamide, 4-chloro-D-phenylalanyl-L-cysteinyl-L-tyrosyl-D-tryptophyl-L-lysyl-L-valyl-L-cysteinyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 84 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1985:572200 HCAPLUS
 DOCUMENT NUMBER: 103:172200
 TITLE: Radioimmunoassay of cholecystokinin: comparison of
 different tracers
 AUTHOR(S): Cantor, Per; Rehfeld, Jens F.

09890219

CORPORATE SOURCE: Dep. Clin. Chem., Univ. Copenhagen, Copenhagen,
DK-2100, Den.
SOURCE: Journal of Immunological Methods (1985), 82(1), 47-55
CODEN: JIMMBG; ISSN: 0022-1759
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The binding of cholecystokinin (CCK) [9011-97-6] antibodies with different sequence-specificities to Bolton-Hunter labeled CCK-33 (125I-BH-CCK-33 [98823-90-6]), CCK-8 (125I-BH-CCK-8 [79672-09-6]) and chloramine-T iodinated gastrin-17 (125I-gastrin-17 [59240-67-4]) was compared. The antibody binding was expressed as the final antiserum dilution (titer) and the effective equilibrium constant of the binding.

Antibodies

specific for the C- or the N-terminal sequence of CCK-8 all bound well to 125I-BH-CCK-8. In contrast, some of the antibodies directed against the common C-terminus of CCK and gastrin displayed remarkably low binding of 125I-gastrin-17 or 125I-BH-CCK-33, whereas all antisera specific for the N-terminal or midsequence of CCK-33 bound 125I-BH-CCK-33 well. The lower binding of 125I-BH-CCK-33 to some C-terminal antibodies raised against gastrin may be due to a C-terminal conformation of CCK-33 different from that of gastrin. In accord with the high specific radioactivity of 125I-BH-CCK-8, the best sensitivity of CCK RIA was obtained with the CCK-8 tracer.

IT 79672-09-6


RL: BIOL (Biological study)
(as tracer, for radioimmunoassay of cholecystokinin peptides)

RN 79672-09-6 HCAPLUS


CN Cholecystokinin-8 (swine), N-[3-[4-hydroxy-3-(iodo-125I)phenyl]-1-oxopropyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



Updated Search



L11 ANSWER 85 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1985:7091 HCAPLUS
 DOCUMENT NUMBER: 102:7091
 TITLE: Renin inhibitors
 INVENTOR(S): Burton, James
 PATENT ASSIGNEE(S): General Hospital Corp., USA
 SOURCE: U.S., 7 pp.
 CODEN: USXXAM
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 4455303	A	19840619	US 1983-497707	19830524
PRIORITY APPLN. INFO.:			US 1983-497707	19830524

OTHER SOURCE(S): MARPAT 102:7091

AB Peptides H-X-X1-X2-X3-X4-R [X, X1, X3 = Phe, Phe(4-Cl), Phe(4-F), Phe(4-Br), Phe(OMe), Tyr, Phe(4-I), Tyr(o-Me); X2 = Val, Thr, threo- α -amino-3-chlorobutyric acid residue; X4 = Lys, Arg; R = NH₂, NHR1 (R1 = C1-4 alkyl), OH, OR2 (R2 = C1-4 alkyl), OM (M = cation)] were prepared as renin inhibitors and they can be used for the treatment of renin-dependent hypertension. Thus, H-Phe(4-Cl)-Phe-Val-Tyr-Lys-NH₂ (I) was prepared by the solid-phase method on a benzhydrylamine resin. I at 70 μ m inhibited human renin by 70%.


IT 91223-94-8P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and renin inhibiting activity of)

RN 91223-94-8 HCAPLUS

CN L-Lysinamide, L-phenylalanyl-4-iodo-L-phenylalanyl-L-valyl-L-tyrosyl-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 86 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1984:552320 HCAPLUS

DOCUMENT NUMBER: 101:152320

TITLE: Synthesis and characterization of an iodinated
derivate of the cholecystokinin octapeptide for
receptor binding studies

AUTHOR(S): Jamieson, James D.; Rosenzweig, Steven A.; Miller,
Laurence J.

CORPORATE SOURCE: Sch. Med., Yale Univ., New Haven, CT, 06510, USA

SOURCE: Biol. Act. Princ. Nat. Prod. (1984), 164-9.

Editor(s): Voelter, Wolfgang; Daves, Doyle G. Thieme:
Stuttgart, Fed. Rep. Ger.

CODEN: 51TMAX

DOCUMENT TYPE: Conference

LANGUAGE: English

AB Cholecystokinin octapeptide (CCK-8), H-Asp-Tyr(SO₃H)-Met-Gly-Trp-Met-Asp-Phe-NH₂, was treated with ¹²⁵I-labeled N-hydroxysuccinimidyl 3-(4-hydroxyphenyl)propionate (¹²⁵I-Bolton-Hunter reagent) in DMF containing Et₃N to give Nα-(¹²⁵I-desaminotyrosyl)-CCK-8 (I). Binding studies of I with rat pancreatic acini showed that interaction with its receptor was rapid, reversible, saturable, and sp. in that only structural analogs of CCK and not unrelated gastrointestinal peptide hormones inhibited binding.



IT 79672-09-6P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and receptor-binding studies of)

RN 79672-09-6 HCAPLUS

CN Cholecystokinin-8 (swine), N-[3-[4-hydroxy-3-(iodo-¹²⁵I)phenyl]-1-oxopropyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 87 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1984:511385 HCAPLUS
 DOCUMENT NUMBER: 101:111385
 TITLE: Analogs of the luteinizing hormone releasing hormone
 having the Azagly10 moiety with antiovolatory activity

AUTHOR(S): Folkers, Karl; Bowers, Cyril Y.; Stepinski, Janusz; Plucinski, Tomasz; Sakagami, Masanori; Kubiak, Teresa

CORPORATE SOURCE: Inst. Biomed. Res., Univ. Texas, Austin, TX, 78712, USA

SOURCE: Zeitschrift fuer Naturforschung, Teil B: Anorganische Chemie, Organische Chemie (1984), 39B(4), 528-32
CODEN: ZNBAD2; ISSN: 0340-5087

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Twenty-four analogs of LH-releasing hormone (LHRH) were synthesized and assayed for antioovulatory activity in rats. [N-Ac-3Δ-Prol, pF-D-Phe2, D-Trp3,6, Azagly10]-LHRH (I, Azagly = NHNHCO) completely inhibited ovulation at 6 µg and was the most potent of the 24; I is a relatively potent antagonist. The Azagly and Ac-NHNH and D-Ala moieties in position 10, and D-Arg in position 6, and diverse substitutions in position 1 were emphasized. D-Arg6 was inferior to D-Trp6, and pCl-D-Phe6 appeared superior to D-Trp6. D-Trp3,6 was superior to D-2-Nal3,6 (D-2-Nal = β-(2-naphthyl-D-alanine residue) and D-His3,6.

IT 91676-08-3P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(preparation and antioovulatory activity of)

RN 91676-08-3 HCAPLUS

CN D-Alaninamide, 1-acetyl-3,4-didehydro-L-prolyl-4-chloro-D-phenylalanyl-4-chloro-D-phenylalanyl-L-seryl-L-tyrosyl-4-chloro-D-phenylalanyl-L-leucyl-L-arginyl-L-prolyl- (9CI) (CA INDEX NAME)

PAGE 1-A




09890219


CN Caerulein, 1-de(5-oxo-L-proline)-2-de-L-glutamine-3-[N-[3-(4-hydroxy-3-iodophenyl)-1-oxopropyl]-L-aspartic acid]-5-L-methionine- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



Updated Search

L11 ANSWER 90 OF 90 HCAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1976:456594 HCAPLUS

DOCUMENT NUMBER: 85:56594

TITLE: Studies on the action mechanism of the antihemostatic effect of iodopeptides

AUTHOR(S): Golub, A. L.; Mende, T. J.

CORPORATE SOURCE: Sch. Med., Univ. Miami, Miami, FL, USA

SOURCE: Thrombosis and Haemostasis (1976), 35(2), 437-46

CODEN: THHADQ; ISSN: 0340-6245

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A synthetic iodopeptide having a glutamic acid-diiiodotyrosine molar ratio of 1:1 (diiiodotyrosine-glutamic acid copolymer [59827-23-5]) was an effective anticoagulant both in vivo and in vitro. Contrasted with heparin [9005-49-6] the following general conclusions may be made regarding its action. The iodopeptide did not act through the inactivation of thrombin in plasma. Iodopeptide did interact with fibrinogen to form a complex which, in vitro, was not soluble in buffered saline at physiol. pH. At pH 8, iodopeptide interacted with fibrinogen to form a soluble complex in the presence of 0.9% NaCl that was not coagulable either by thrombin or Crotalus venom enzymes. All the available evidence indicates that the fibrinogen to fibrin conversion was not inhibited under these conditions, but that fibrin, once formed, was not able to polymerize due to interference by iodopeptide. Similar results were obtained with heparin in vitro with thrombin-fibrinogen mixts. in the absence of NaCl. Studies with Russell's viper venom in native PRP strongly suggest that the iodopeptide also interferes with processes in the early coagulation pathway associated with prothrombin activation.

IT 59884-15-0

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(anticoagulant activity of)

RN 59884-15-0 HCAPLUS

CN L-Glutamic acid, N-[N-[N-[N-[N-[N-[N-(N-L- α -glutamyl-3,5-diiodo-L-tyrosyl)-3,5-diiodo-L-tyrosyl]-L- α -glutamyl]-3,5-diiodo-L-tyrosyl]-L- α -glutamyl]-3,5-diiodo-L-tyrosyl]-L- α -glutamyl]-3,5-diiodo-L-tyrosyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

